

The Limits to Stock Return Predictability

Donald Robertson and Stephen Wright*

April 2, 2009

Abstract

We examine predictive return regressions from a new angle. We ask what observable univariate properties of returns tell us about the “predictive space” that defines the predictive model: the triplet (λ, R_x^2, ρ) , where λ is the predictor’s persistence, R_x^2 is the predictive R-squared, and ρ is the "Stambaugh Correlation" (between innovations in the predictive system). When returns are nearly white noise and the variance ratio of long-horizon returns slopes downwards we show that the predictive space can be quite tightly constrained. Data on real annual US stock returns suggest there is a very limited scope for even the best possible predictive regression to out-predict the univariate representation, particularly over long horizons.

Keywords: Predictive return regressions, variance ratio, fundamental and non-fundamental univariate representations.

*Faculty of Economics, University of Cambridge, dr10011@econ.cam.ac.uk and Dept of Economics, Birkbeck College, University of London, s.wright@bbk.ac.uk

1 Introduction

A perennial problem in empirical finance is the question of whether it is possible to predict stock market returns using some predictor variable. In addition to the question of what predictor variable to use (the list of potential predictors includes, *inter alia*, dividend yields, interest rates, book-market values, price-earnings ratios), there are econometric questions regarding the appropriate distributional theory for inference (Stambaugh 1999, Campbell and Yogo 2006, Ang and Bekaert 2006 and many others); whether any underlying relationships are stable enough to allow useful predictability; or may simply arise from data mining (Pesaran and Timmerman 1995, Timmermann and Paye 2006, Ferson et al 2003, Campbell and Thomson 2008, Cochrane 2008, Goyal and Welch 2003). Finally there is a closely related literature that addresses differences between one period ahead and long horizon regressions (see for example Campbell and Viceira 2002; Cochrane, 2008; Boudoukh et al, 2008).

In this paper we examine predictive return regressions from a new angle. It is well-known that when one time series predicts another the properties of the predictive system (including the univariate properties of the predictor variable) determine the univariate properties of the predicted variable (in this case returns). But we can also view the process in reverse. In this paper we ask what the observable univariate properties of returns tell us about the “predictive space” that defines the predictive model: the triplet (λ, R_x^2, ρ) where λ is the predictor’s persistence, R_x^2 is the predictive R-squared, and ρ is the "Stambaugh Correlation" (between the innovations to the predictor autoregression and those in the predictive return regression).¹

If these insights depended on tight estimates of parameters in an ARMA representation our analysis would not be of much practical value. But in fact the reverse is the case. The very fact that ARMA representations of returns fit so *poorly* is informative about the nature of the predictive space for predictive models in general. But a second feature of returns data also has potentially very significant implications for the predictive space: whether the variance ratio of long-horizon returns slopes downwards. The academic literature that directly tests for this latter feature has not yielded unanimous conclusions (contrast, for example, the original

¹In this paper we focus on a widely used predictive regression framework that can be reduced to just three parameters. Some of our key results also extend to more general predictive models.

evidence presented by Poterba and Summers 1988 with the revisionist approach of Kim et al 1991 and Pastor & Stambaugh, 2008). But we would argue that the implications of “variance compression”² are worth considering, first, because it is usually taken for granted (whether explicitly or implicitly) by investment practitioners as the basis for the buy-and-hold strategy (for the classic explicit statement of this view see Siegel, 1998); and second, because the analysis we present in this paper will lead us to argue that it is also *implicit* in a much wider range of literature that assumes predictability of returns, especially over long horizons (for example, Campbell & Viceira 1999, Cochrane, 2008).

Since a declining variance ratio is a univariate property it cannot co-exist with returns being *completely* unpredictable from their own past, although we show that there is no necessary contradiction between a quite significantly declining variance ratio at long investor horizons and a very weak degree of short-term univariate predictability. However, the combination of these two features *does* have significant implications for the predictive space that contains all possible predictive regressions. We show that the predictive space for stock returns can quite easily contract to such an extent that there is little, if any, scope for predictive regressions to out-predict the univariate representation, particularly at long horizons.

Recent stock market movements have been a reminder of the continuing significance of this issue. Many predictors of stock returns originate as valuation indicators. In the late 1990s most were signalling that the market was “over-valued”, in the very broad sense of the word³ that such indicators were predicting weak returns (see, for example, Campbell & Shiller, 1998; Fama & French, 2002; Shiller, 2000). In the more recent past, as markets have weakened sharply, the issue has arisen of when they become sufficiently “cheap” to offer a good prospect of unusually strong returns. Most such indicators can be interpreted as the ratio of the stock price to some underlying fundamental. Our analysis suggests that even if the best possible predictor were found for the stock market (which could always be expressed as a ratio of the stock price to some best possible measure of fundamentals) it might barely predict any better than simply

²For reasons noted below we prefer this term to the more commonly used “mean reversion”, which we argue is a misnomer.

³That is, without putting any necessary interpretation in terms of market efficiency: contrast the perspectives of Shiller, 2000 and Fama & French, 2002, for example.

using the history of returns to forecast future returns.

The paper proceeds as follows.

We first, in Section 2, set out the evidence for the two key empirical features of stock returns that motivate our analysis: a low univariate R-squared and a declining variance ratio. We then show, in Section 3, how the standard predictive regression framework determines the univariate representation of returns, which is an ARMA(1,1).

Our key results are in Section 4, where we show how the univariate properties of returns can restrict the predictive space. We show that we can derive lower and upper bounds for the predictive R-squared, R_x^2 , that depend solely on univariate properties. We then show that a declining variance ratio for long-horizon returns implies that the "Stambaugh Correlation, ρ , is in general bounded away from zero, and for a plausible range of ARMA parameters can be very close to unity. A high Stambaugh Correlation is usually treated as a nuisance that complicates inference; we show that it is an intrinsic characteristic of the true predictor of stock returns if there is variance compression. Finally we show a further feature that arises from these restrictions on the predictive space: the correlation between the true predictor and a "pseudo predictor", derived solely from the history of returns, is also bounded away from zero. It approaches unity as the predictive space contracts; however even when the predictive space is quite tightly constrained this correlation can still be a reasonably long way below unity.

In Section 5 we provide numerical and empirical illustrations. In the benchmark case of pure white noise returns, the higher is λ , the AR(1) parameter of the predictor, the more constrained is the predictive space. But for given λ , any degree of variance compression must cause the predictive space to contract relative to the white noise benchmark. Point estimates derived from estimated ARMA representations of returns, suggest that the predictive space is very tightly constrained. The estimate of the best possible predictive R-squared of any possible predictor is around 10% and that of the minimum Stambaugh Correlation is at least 0.9. The implied lower bound for the correlation between the best possible predictor and the "pseudo predictor" is lower, but still a long way from zero. These estimates therefore suggest limited scope for even the best possible predictor of returns to out-perform the univariate representation.

Given the imprecision of ARMA estimation, we acknowledge that a quite wide range of

near-white noise processes could also be generating the return series, for some of which the predictive space is less tightly constrained. However, a key requirement for a significantly *less* constrained predictive space is that the true predictor must have quite low persistence. Our analysis therefore suggests two simple pre-tests for potential predictors of stock returns: they should not look too similar to the “pseudo predictor” that summarises the history of returns; and they should not be too persistent. We note that commonly used predictors of stock returns do not match up to either of these criteria.

Finally, in Section 6 we note a further implication of our analysis for the the predictability literature. The strength of long-horizon return predictability is driven by a combination of predictor persistence and a declining variance ratio; but it is exactly these features that lead to a tightly restricted predictive space. We therefore conclude that long-horizon return predictability, if it exists, must be close to being a univariate phenomenon.

In Section 7 we draw conclusions and implications of our analysis; appendices provide algebraic derivations and proofs.

2 Univariate features of returns


As a preamble to our analysis we briefly summarise the observable features of returns that we shall draw on in the rest of our analysis.

2.1 Returns are near-white noise

In Figure 1 we plot the autocorrelation function for real annual stock returns in the United States over two samples, 1871-2008 and 1945-2008. The first spans the full available dataset on a reasonably consistent basis for a broad based US stock market measure (the Cowles (1938) industrial index from 1871-1925, and the S&P 500 thereafter).⁴ The shorter sample allows for the possibility that return properties may have changed in the postwar era (consistent with the claims discussed below by Kim et al, 1991).

⁴We have also extended this series backwards to 1801 using Siegel’s (1998) dataset. Results are very similar.

Figure 1. The Autocorrelation Function of Real US Stock Returns



We also show bootstrapped 5% and 95% bounds when returns are resampled with replacement to destroy any possible temporal dependence. In the full sample this illustrates that, whilst autocorrelations are generally very small in absolute terms, a subset are individually marginally significant against the null of white noise; the same applies for the standard Ljung-Box Q portmanteau test. However even these apparent rejections of the white noise null are subject to a well-known data mining critique, if we focus only on a relatively small number of rejections. In Table 1 we show simulated p-values for the largest absolute autocorrelation over a different range of lag lengths up to some maximum, over the two different samples, and for the most significant rejection on the Ljung-Box test, both under the null of white noise. This shows that even white noise processes will appear to have significant autocorrelations at *some* lag length with quite high probability; with the probability increasing with the total number of autocorrelations considered. Thus on the basis of standard analysis of autocorrelations, returns appear to be very close to white noise even over the full sample. In the postwar sample, there is even less reason to reject the white noise null.

Of course, as is equally well-known, tests of the white noise null will have very low power against an alternative that the true process is close to, but is not quite white noise. But for our purposes, it turns out that the distinction is not of any great importance. We shall show below that even if we allow returns to deviate from white noise by estimating ARMA(1,1)

representations (which appears to be quite adequate to remove any serial correlation structure in the resulting residuals) the resulting representations have very low R-squareds.

Thus the first key feature that informs our analysis is that returns are, at best, barely predictable in terms of their own past.

| Table 1. Bootstrapped p-values under the white noise null⁵ | | | | | | |
|---|-------------------------------|-----------|-------------------------------|-----------|----------------------|-----------|
| | max(Absolute Autocorrelation) | | min (p -value on Q test) | | min (Variance Ratio) | |
| h_{\max} | 1871-2008 | 1945-2008 | 1871-2008 | 1945-2008 | 1871-2008 | 1945-2008 |
| 10 | 0.089 | 0.797 | 0.095 | 0.643 | 0.225 | 0.648 |
| 20 | 0.196 | 0.835 | 0.142 | 0.762 | 0.165 | 0.761 |
| 30 | 0.314 | 0.967 | 0.193 | 0.796 | 0.032 | 0.155 |
| 40 | 0.452 | 0.973 | 0.262 | 0.554 | 0.074 | 0.030 |

2.2 The variance ratio slopes downwards

In Figure 2 we plot the variance ratio for real annual stock returns at horizon h , $VR(h) = Var(\sum_{i=1}^h r_{t+i}) / (Var(r_t).h)$ for horizons 1 to 40.⁶

The first panel shows that over the long sample 1871-2008 the variance ratio shows clearly the pattern identified by Poterba & Summers, 1988. It declines nearly monotonically as the horizon increases until around $h = 30$, at which point it appears to level out at a value of around 0.2: thus indicating a reduction in volatility for long-horizon returns that is, in economic terms, highly significant, since it implies that long-term investment in stock portfolios has been very much less risky than it would have been if returns were white noise.⁷ We also show simulated


⁵We simulate the white noise null by resampling with replacement from the empirical distribution of real annual stock returns, in 10,000 repetitions. The first two columns of Table 1 show the bootstrapped probability of a larger value than in the data for the maximum autocorrelation from 1 to h_{\max} , under the white noise null. For Columns 3 and 4, we carry out Ljung-Box Q tests of the joint significance of autocorrelations from 1 to h , in the data, and in each replication, then find the minimum nominal p -value over $h=1$ to h_{\max} : the table shows the probability, across all replications, of observing a lower minimum nominal p -value than in the data. In Columns 5 and 6 we show the bootstrapped probability across all replications, of a lower minimum variance ratio than in the data over horizons 1 to h_{\max} .

⁶We do *not* include the small sample adjustment proposed by Cochrane (1988) and others. Given our focus on simulated results, where the variance ratio is calculated in the same way in both data and simulations, any adjustment is unnecessary. Under the white noise null the unadjusted sample variance ratio is biased downwards; however under alternatives where returns are near-white noise such as the ARMA(1,1) we analyse below, we show that the unadjusted ratio appears to be close to being unbiased.

⁷This is the statistical basis for the dominant investment strategy of buy-and-hold. The classic argument on this basis is in Siegel, 1998. Campbell & Viceira (1999) discuss the implicit contradictions in the logic of the

5% and 95% bounds for the sample variance ratio under the bootstrapped white noise null. The observed pattern does not differ much from white noise at short horizons; but appears increasingly different as the horizon lengthens. While the data mining critique again argues against placing too much weight on individual horizons, the third and fourth column of Table 1 shows that if we focus on the minimum variance ratio across all horizons up to a given maximum horizon, the longer the horizon, the lower is the probability of observing such a low value under the white noise null.

Figure 2. The Variance Ratio of Real US Stock Returns



The second panel of Figure 2 shows that if we calculate the variance ratio only over the postwar period there is no systematic tendency to decline until $h = 20$ - a result consistent with the estimates in Kim et al (1991). However, for longer horizons the decline is quite marked, and, as Table 1 shows, statistically significant even allowing for the very limited number of degrees of freedom.⁸ We shall also show below that there are clear *indirect* measures of a declining variance ratio that persist into the postwar era.

As we noted in the introduction, if the variance ratio for stock returns is declining, this cannot be consistent with returns being white noise, since, as Cochrane's (1988) original analysis strategy, which advocates that investors should pay attention to the benefits of the (weak) implied predictability of returns, but not actually exploit this predictability by changes in weightings on stocks.

⁸The downward bias noted in footnote 6, which is quite severe in such a relatively short sample, is very evident in the simulated upper and lower bounds.

showed, we have $VR(h) = 1 + 2 \sum_{j=1}^{h-1} \left(\frac{h-j}{h}\right) corr(r_t, r_{t-h})$. But there is no necessary contradiction between our weak results for the autocorrelation function and the stronger results for the variance ratio, since the latter relates to a long weighted average of autocorrelations. In principle “variance compression”⁹ can be both quite significant, and consistent with a very limited degree of short-term predictability. This is indeed what appears to be the case in the data.

It should also be stressed that the probability that *both* these features would appear in the data would be *very* small under the white noise null. Figure 3 illustrates for the full sample. We resample 138 observations of the real stock return to simulate the white noise null. Figure 3 is then a scatter plot of the minimum variance ratio, over horizons 1 to 40, against the sample R-squared for an ARMA(1,1) representation of returns, for each replication. The crossing point of the two lines on the chart shows the values observed in the data.

Figure 3 shows that the bulk of replications would have low ARMA R-squareds, but with considerable spread: 24% of the simulated values would be above the value in the data (very much in line with the evidence on the autocorrelations shown in Table 1). The majority of simulations would generate a minimum variance ratio well above that in the data; the points below the horizontal line correspond to the 7.4% probability given in the bottom row of Table 1. But, most strikingly, samples in which the variance ratio does appear to slope significantly downwards are almost always also samples in which the ARMA model appears to predict distinctly better than in the data: only 1.4% of replications generated combinations in the bottom left quadrant, ie, with both a lower R-squared *and* a lower minimum variance ratio than in the data.¹⁰


⁹This feature is often referred to as mean reversion (following Poterba & Summers, (1988)), but this is a somewhat confusing misnomer. Poterba & Summers define mean reversion as "stock prices (or cumulative returns) have a mean-reverting transitory component". Following Beveridge & Nelson (1981) we can write any general ARMA(p, q) univariate representation of returns as

$$r_t = a(L)\varepsilon_t = a(1)\varepsilon_t + a^*(L)(1-L)\varepsilon_t$$

with the second term defining the mean-reverting transitory component in cumulative returns $= a^*(L)\varepsilon_t$. Such a term will be present for *any* stationary univariate representation where returns have some serial correlation structure, but not all such representations will have a downward sloping variance ratio. It is straightforward to show that $a(1) < 1$ is a sufficient condition for the variance ratio to slope downwards. Since $a(1) + a^*(0) = 1$, in this case the transitory component will be positively correlated with returns, whereas for $a(1) > 1$, which implies that the variance ratio slopes upwards, it will be negatively correlated. But in both cases the transitory component will be mean-reverting (cf Kim et al (1991) who refer to the latter case as “mean aversion”).

¹⁰An even lower percentage (1.0%) lie in this quadrant in the postwar sample.

Figure 3. Univariate Predictability and a Declining Variance Ratio:



3 The predictive regression framework

3.1 The general system

Consider the system used by Stambaugh (1999) and many others in the analysis of predictive return regressions

$$r_t = -\beta_x x_{t-1} + u_t \tag{1}$$

$$x_t = \lambda x_{t-1} + v_t \tag{2}$$

where the first equation captures the degree of predictability of some variable r_t , typically stock returns or excess returns over some interval, in terms of a predictor variable x_{t-1} , and the second describes the autocorrelation of the predictor variable. We assume $0 \leq \lambda < 1$; $\beta_x \geq 0$ so that both r_t and x_t are stationary.¹¹ We put no restrictions on the innovations u_t and v_t other than that they be (jointly) serially uncorrelated mean zero with finite variance. We assume all data are zero mean for simplicity, hence neglect constants.

Equation (1) is quite general, since x_{t-1} may in principle be some weighting of a set of

¹¹We can always change the sign of x to ensure $\beta_x \geq 0$. The restriction that β_x be non-negative simplifies subsequent algebra. Most of our results generalise to, but are complicated by, $\lambda < 0$; however, given our discussion above relating to *a priori* properties of predictors we regard this as empirically less likely to be of interest.

variables with predictive power for r_t and the error term may capture a range of nonlinearities. Equation (2) is distinctly more restrictive, but, since Stambaugh has been widely used in the literature and, again, allowing for exotic errors, can still encompass a wide range of models (including for example two state Markov switching models, Hamilton 1989).

Substituting from (2) into (1) we derive the reduced form process for r_t , which is an ARMA(1,1):

$$r_t = \lambda r_{t-1} + \varepsilon_t - \theta \varepsilon_{t-1} = \left(\frac{1 - \theta L}{1 - \lambda L} \right) \varepsilon_t \quad (3)$$

where L is the lag operator, such that $Lx_t = x_{t-1}$; ε_t is a serially uncorrelated innovation; and as long as u_t and v_t are less than perfectly correlated, we can choose the “fundamental” solution for the MA parameter that has $\theta \in (-1, 1)$, so the representation is strictly invertible in terms of the history of r_t .¹² If $\theta = \lambda$ the AR and MA components cancel, and r_t will be white noise (note that this does *not* require $\beta_x = 0$).

We first note that in the ARMA representation the properties of r_t are entirely determined, up to a scaling factor, by the pair (λ, θ) . The properties of the underlying predictive system (1) and (2) can in turn be characterised by the three unit-free parameters (λ, ρ, R_x^2) where $\rho = \sigma_{uv}/(\sigma_u \sigma_v)$ is the correlation between the innovations in the model, and $R_x^2 = 1 - \sigma_u^2/\sigma_r^2$ is the R^2 in the predictive regression. We shall refer to the triplet (λ, ρ, R_x^2) as the “predictive space”.

The autoregressive coefficient of the predictor variable translates directly to the AR coefficient of the reduced form (3). For the case of the MA parameter θ things are more complicated. In Appendix A we show that θ depends on all three parameters that define the predictive space,

$$\theta = \theta(\lambda, \rho, R_x^2) \quad (4)$$

We shall show that the two univariate properties summarised Section 2 mean that the predictive space can be quite tightly constrained. In so doing it will be helpful to make reference to two important benchmark cases that we shall show determine the nature of these restrictions.

¹²See Appendix A.

3.2 “Pseudo Predictor” Representations

In this section we define two special cases of the predictive system in (1) and (2), both of which can be derived directly from the properties of the ARMA representation. We shall then go on to show, in Section 4, that these special cases provide benchmarks that allow us to set limits on the “predictive space” that contains *all* possible predictive systems of the form in (1) and (2).

3.2.1 The fundamental pseudo predictor

If we start from the ARMA(1,1) representation in (3) we can also reparameterise it as a predictive system of the same general form as (1) and (2):¹³

$$r_t = -\beta_f x_{t-1}^f + \varepsilon_t \quad (5)$$

$$x_t^f = \lambda x_{t-1}^f + \rho_f \varepsilon_t \quad (6)$$

where we define the “fundamental pseudo predictor” by

$$x_t^f = \text{sign}(\theta - \lambda) \frac{\varepsilon_t}{1 - \lambda L} \quad (7)$$

and set $\beta_f = |\theta - \lambda|$ and $\rho_f = \text{sign}(\theta - \lambda)$. The definition in (7) preserves the useful property that β_f , like β_x , is always positive.

The fundamental pseudo predictor has the same AR(1) form as the true predictor variable, but will generate identical predictions to the fundamental ARMA representation in (3). It will therefore have the same predictive R-squared as the ARMA, which we show in Appendix B is given by

$$R_f^2(\lambda, \theta) \equiv 1 - \frac{\sigma_\varepsilon^2}{\sigma_r^2} = \frac{(\theta - \lambda)^2}{1 - \lambda^2 + (\theta - \lambda)^2} \quad (8)$$

¹³See Appendix A for derivation.

Note that, using (7) and (3) we can also write

$$x_t^f = \text{sign}(\theta - \lambda) \frac{r_t}{1 - \theta L} = \text{sign}(\theta - \lambda) \sum_{i=0}^{\infty} \theta^i r_{t-i} \quad (9)$$

so the fundamental pseudo predictor is simply a geometrically weighted average of lags of r_t where the weights are given by powers of θ .

3.2.2 The non-fundamental pseudo predictor

For every fundamental ARMA(1,1) representation there is an associated “non-fundamental” representation, given by

$$r_t = \left(\frac{1 - \theta^{-1}L}{1 - \lambda L} \right) \eta_t \quad (10)$$

with $\sigma_\eta^2 = \theta^2 \sigma_\varepsilon^2$. This representation generates an identical autocorrelation structure for returns to the fundamental representation, but, as is well known (see for example Hamilton, 1994, pp 66-67), the non-fundamental innovations, η_t cannot be recovered from the history of r_t , hence the non-fundamental representation does not represent a viable predictive model. As a result, with a few exceptions (most notably, Lippi & Reichlin 1994; Fernandez-Villaverde et al, 2007) non-fundamental representations have received relatively little attention.

To see why η_t cannot be recovered from the data, note that if we attempt to solve (10) for η_t we have

$$\eta_t = \left(\frac{1 - \lambda L}{1 - \theta^{-1}L} \right) r_t = \sum_{i=0}^{\infty} \theta^{-i} [r_{t-i} - \lambda r_{t-i-1}]$$

given that $|\theta^{-1}| > 1$ the sum does not converge, hence the representation in (10) is not invertible in terms of the history of r_t . However, if (strictly hypothetically) we had data on current and *future* values of r_t , we could write

$$\eta_t = \left(\frac{1 - \lambda L}{1 - \theta^{-1}L} \right) r_t = -\theta F \left(\frac{1 - \lambda L}{1 - \theta F} \right) r_t = -\theta \sum_{i=1}^{\infty} \theta^i [r_{t+i} - \lambda r_{t+i-1}] \quad (11)$$

where F is the forward shift operator, such that $Fx_t = L^{-1}x_t = x_{t+1}$, and in this case the sum does converge. Thus the non-fundamental ARMA representation *does* have an invertible

representation, but only in terms of current and future values of r_t , making it valueless as a predictive model.

Of course, if we already knew the entire future of r_t , we would not need a predictive model at all, therefore there would be no point in constructing a series for η_t . But the reverse is not the case. In general, even if we did have data on η_t (perhaps by some divine dispensation) this would *not* reveal the entire future of r_t , but rather a particular linear combination of future values. Thus while the non-fundamental representation would, if we had the history of η_t , predict better than the fundamental representation, it would not in general predict perfectly.¹⁴

While it may seem somewhat peculiar to take an interest in a predictive model that is so manifestly non-viable, it turns out that it provides us with an extremely useful benchmark. And it does so because, while we will never, in practice, be able to observe η_t , we do know the predictive *properties* of the non-fundamental representation, even if we cannot actually use it to predict, since these can be inferred directly from the properties of the fundamental representation.¹⁵

Thus, as noted above, the equivalence of the two representations must imply that η_t has lower variance than ε_t , the fundamental innovation ($\sigma_\eta^2 = \theta^2 \sigma_\varepsilon^2$), hence, if we did have data on η_t , the non-fundamental representation would predict strictly better than the fundamental representation. Its (strictly notional) predictive R-squared can be derived by replacing θ with θ^{-1} in (8), giving

$$R_n^2(\lambda, \theta) \equiv 1 - \frac{\theta^2 \sigma_\varepsilon^2}{\sigma_r^2} = \frac{(1 - \theta\lambda)^2}{1 - \lambda^2 + (\theta - \lambda)^2} > R_f^2(\lambda, \theta) \quad (12)$$

We can also, as for the fundamental ARMA representation, again reverse-engineer a representation of the same general form as (1) and (2), ie if we define a "non-fundamental pseudo

¹⁴Except in the limit as $\theta \rightarrow 0$. See Appendix D.

¹⁵Strictly speaking in Lippi & Reichlin's (1994) terminology the non-fundamental representation in (10) is a "basic" non-fundamental representation, in that it is of the same order as the observable fundamental representation. There is in principle an infinity of "non-basic" non-fundamental representations of arbitrary higher order, since any white noise innovation can always be given a non-fundamental representation: ie, we could write $\eta_t = (1 - \phi^{-1}L)(1 - \phi L)^{-1} \omega_t$, with $\sigma_\omega^2 = \phi^2 \sigma_\eta^2$, and in principle then find a non-fundamental representation of ω_t , and so on ad infinitum. But nothing in the data tells us anything about ϕ , and hence about ω_t , hence we can infer nothing from the data about the properties of such non-basic representations.

predictor", by

$$x_t^n = \text{sign}(\theta^{-1} - \lambda) \frac{\eta_t}{1 - \lambda L} \quad (13)$$

we can write

$$r_t = -\beta_n x_{t-1}^n + \eta_t \quad (14)$$

$$x_t^n = \lambda x_{t-1}^n + \rho_n \eta_t \quad (15)$$

with $\beta_n = |\theta^{-1} - \lambda|$ and $\rho_n = \text{sign}(\theta^{-1} - \lambda)$.

3.3 The variance ratio in the ARMA(1,1) reduced form

We have already referred, in our discussion of univariate properties in the data, to the variance ratio at horizon h , as originally defined for the general case by Cochrane (1988) as

$$VR(h) = \frac{1}{h} \frac{\text{Var}(\sum_{i=1}^h r_{t+i})}{\text{Var}(r_t)} \quad (16)$$

It is straightforward to show¹⁶ that, for the ARMA(1,1) process (3), that is the reduced form of the predictive system, $VR(h)$ is monotonic in h and that

$$VR(h) \begin{cases} < 1 \Leftrightarrow \theta > \lambda \\ > 1 \Leftrightarrow \theta < \lambda \end{cases} ; \forall h > 1 \quad (17)$$

We shall also make use of the limiting value of the variance ratio, which in the ARMA(1,1) can be expressed as

$$V = \lim_{h \rightarrow \infty} VR(h) = (1 - R_f^2) \left(\frac{1 - \theta}{1 - \lambda} \right)^2 \quad (18)$$

where, given the monotonicity of the $VR(h)$ in h , we also have $V < 1 \Leftrightarrow VR(h) < 1 \forall h > 1$.

¹⁶See Appendix C.

4 The Predictive Space for Stock Returns

4.1 Bounds on the ARMA(1,1) coefficients, θ and λ



In Section 2 we discussed two univariate properties of real stock returns in the data: first that they were near-white noise (hence barely predictable in the short term); and second that there appears to be quite strong evidence of “variance compression” (ie, a declining variance ratio). It is straightforward to show that these two features of the data constrain quite tightly the possible values of λ and θ in the ARMA representation. We shall subsequently see that this in turn will place quite significant restrictions on the predictive space.

Using the ARMA(1,1) framework outlined above we plot, in Figure 4, contours in (θ, λ) space of equal R_f^2 and of equal V (the limiting value of the variance ratio).¹⁷ The top panel of Figure 4 shows contours for $V = 0.4$ and $R_f^2 = 0.05$. The shaded area then gives the admissible set of (θ, λ) that generate values of V no greater than 0.4 and R_f^2 of no more than 0.05. Lower values of V push the V -contour up and to the left, while lower values of R_f^2 move the R_f^2 -contours towards the 45 degree line, thus reducing the admissible (θ, λ) area. The second panel of Figure 4 illustrates: the shaded area is now the (θ, λ) combinations consistent with V no greater than 0.2, and R_f^2 below 0.025: the permissible space for both ARMA parameters becomes very tightly constrained.¹⁸ Thus if stock returns have little one period ahead univariate predictability, and declining variance ratios, this requires that λ be quite large and that ρ and R_x^2 be such that $\theta(\lambda, \rho, R_x^2) > \lambda$.

¹⁷For given values of R_f^2 and V , we solve (8) and (18) for θ in terms of λ . The former gives two solutions for θ , symmetric around the 45 degree line.

¹⁸The numbers used for R_f^2 and V in Figure 4 are illustrative, but are quite consistent with the evidence illustrated in Figures 2 and 3. Sample estimates of R_f^2 are, if λ is high, subject to severe Stambaugh (1999) bias. Simulation evidence shows that even in a sample as long as the 1871-2008 period discussed in Section 2 a true R_f^2 of 0.025 would result in a mean sample estimate at least twice as large, thus consistent with what we observe in the data. For large λ and θ we can also have $VR(h)$ well above the limiting value V even for horizons as long as those shown in Figure 2.

Figure 4
The Permissible Space for ARMA Parameters for Stock Returns



We showed in Section 3 that the ARMA representations inherits the AR parameter λ from the true predictor. While Figure 4 shows that the requirement that λ be large arises naturally from the univariate properties of returns, we would also argue for this feature on *a priori* grounds. Persistence of a very wide range of economic time series is arguably endemic (if it were not then the massive econometric literature on unit and near-unit root processes would never have arisen). As we noted in the introduction, both observable and notional unobserved predictors of returns can always be conceptualised as the ratio of the stock price to some (usually slow-moving) measure of the “fundamental”. In the long term if the predictor is stationary, equilibrium price and fundamental must move together - ie, must be cointegrated.¹⁹ It is well documented that long-run stationary relations often have very slow adjustment speeds. Virtually all observable predictors of stock prices (most notably valuation ratios like the price-dividend ratio or the price-earnings ratio) have this characteristic.²⁰ But the analysis illustrated by Figure 4 shows that, for sufficiently strong variance compression, and sufficiently weak short-term univariate predictability, the same must apply for *any* logically possible predictor, and that *a priori* arguments and the properties of stock returns are thus mutually consistent.

We now go on to show that these univariate features can put significant restrictions on the predictive space of the underlying model that generates them.

4.2 Bounds for the one-period-ahead predictive R^2

Proposition 1 *For a fundamental ARMA(1,1) representation of returns which is a reduced form of a predictive regression (1) and a predictor autoregression (2) with less than perfectly correlated innovations ($\rho \in (-1, 1)$) the one-period-ahead R^2 of the predictive regression, R_x^2 , satisfies the inequality*

$$R_f^2(\lambda, \theta) < R_x^2 < R_n^2(\lambda, \theta) \tag{19}$$

where R_f^2 and R_n^2 are as defined in (8) and (12).

¹⁹For a powerful exposition of the implications of this form of relationship, see Cochrane, 2008.

²⁰The AR(1) coefficients for the dividend yield and the cyclically adjusted P/E multiple, for example, are 0.92, 0.93.

Proof. See Appendix D.²¹ ■

The lower bound for R_x^2 is the predictive R-squared of the fundamental ARMA representation, or, equivalently, of the “fundamental pseudo predictor” defined in Section 3.2.1. As such it is quite easy to interpret. The true predictor provides predictive information for r_t beyond that contained in the history of r_t itself; it must therefore have a higher predictive R-squared.²²

The upper bound for R_x^2 is the predictive R-squared of the *non*-fundamental ARMA representation, or equivalently of its associated pseudo predictor, defined in Section 3.2.2. The basic intuition for this result can be related to our earlier discussion of the properties of the non-fundamental representation. We showed in Section 3.2.2 that the non-fundamental innovation η_t can be expressed, in (11), as a linear combination of current future returns: so we know already that it must have some predictive power beyond that already in the history of returns. But the result in Proposition 1 is stronger: it shows that the non-fundamental pseudo predictor in period t is the best *possible* predictor of r_{t+1} consistent with its observable univariate properties.²³

We shall show that when the ARMA parameters, θ and λ , lie within the permissible range illustrated in Figure 4, for a given degree of variance compression and low short-term predictability then the allowable range of R_x^2 given by Proposition 1 can become quite small.

Our focus thus far has been on just two of the elements in the predictive space, namely λ and R_x^2 . But a further important feature of Proposition 1 is that both the upper and lower bounds arise in limiting cases of the predictive system in which the Stambaugh Correlation, ρ , is precisely unity in absolute value.²⁴ We now examine intermediate cases in which the innovations are not perfectly correlated.

²¹In Robertson & Wright 2009a we show that this inequality can be generalised to a general ARMA(p, q), but the proof is more convoluted, and the ARMA(1,1) considered here has the dual advantage that it can be related much more readily to the predictive regression framework, and yields relatively simple analytical results.

²²An alternative interpretation of the fundamental pseudo predictor is as the optimal estimate of the true predictor given the information set $\{y_i\}_{i=-\infty}^t$, using the Kalman Filter. If we have data on the true predictor, rather than its estimate, we must be able to predict better.

²³Given the ARMA(1,1) property of returns we know that the true predictor must be an AR(1). For reasons discussed in Footnote 16 there may in principle be multiple predictors, of arbitrary ARMA order, that predict arbitrarily better. But the data tell us absolutely nothing about such predictors. Equivalently, the non-fundamental ARMA representations associated with such predictors are, in Lippi & Reichlin’s (1994) terminology, “non-basic”.

²⁴This is evident from the specification of the two limiting systems, in (5) and (6), and in (14) and (15).

4.3 Bounds for ρ for predictor variables

We have from (4) that the ARMA coefficients θ and λ are linked to the predictive R_x^2 and the Stambaugh Correlation between the innovations, ρ , by $\theta = \theta(\lambda, \rho, R_x^2)$. Thus for a given (θ, λ) pair there is a contour of possible consistent values of (R_x^2, ρ) . Again it turns out that the univariate properties of r_t impose limits on the possible values of ρ .

Proposition 2 *Consider a fundamental ARMA(1, 1) univariate representation (3) which is a reduced form of a predictive regression (1) and a predictor autoregression (2). For $0 < \lambda < \theta$ (ie the variance ratio slopes downwards and the predictor has positive persistence) the correlation coefficient ρ between the two disturbances has a minimum value ρ_{min} which is strictly positive.*

Proof. See Appendix E. ■

Figure 5 illustrates. We graph the contours in (R_x^2, ρ) space for a range of different representations of the return process.

As a benchmark, the lowest contour line shows combinations of the two parameters consistent with a strictly white noise process when the predictor is quite strongly persistent ($\theta = \lambda = 0.78$). The better the predictive model, the higher the associated Stambaugh Correlation, ρ , must be. While ρ can take any value in $[0, 1]$, the lower bound for ρ will only be attained with $R_x^2 = 0$. Thus even in the white noise case a *useful* predictor must have positive ρ .²⁵

The remaining contour lines represent a range of near-white noise processes, all with the same univariate R-squared ($R_f^2 = 0.025$) but with progressively stronger degrees of variance compression (ie, lower values of V). For a given degree of short-term predictability, this corresponds to a progressive reduction in the upper bound for R_x^2 . Since $\rho = 1$ at both upper and lower bounds, the range of possible values of ρ is progressively reduced, hence ρ_{min} in Proposition 2 becomes closer and closer to unity.

²⁵For the white noise case R_f^2 is of course always zero, and the formula for R_n^2 in (36) simplifies to $R_n^2(\lambda, \lambda) = 1 - \lambda^2$, hence, since $\max(R_x^2) = R_n^2(\lambda, \lambda)$ as λ increases the contour line rotates anti-clockwise.

Figure 5. The Predictive Space for Stock Returns: R_x^2 and ρ





Figure 6. The correlation between the true predictor and the fundamental pseudo predictor



This feature of our results sheds light on a significant feature of the the empirical literature on predictive regressions. In this literature a high value of the Stambaugh Correlation is usually treated simply as a nuisance that complicates inference. Our results show that when returns have declining variance ratios (or even if they are purely white noise) it is an intrinsic feature

of the true predictor of returns.²⁶

4.4 How different are predictor variables from the history of returns?

We have shown in the previous section that the Stambaugh Correlation is likely to be close to unity (ie, innovations to the predictor variable will be strongly correlated with innovations in the predictive regression). We also know that, by construction, the fundamental pseudo predictor, which from (9) is simply a weighted average of past returns, has a Stambaugh Correlation of precisely unity. It might therefore seem that any predictor must resemble the pseudo predictor quite closely. In fact, while this may be the case for certain univariate processes, the correlation between the true predictor and the fundamental pseudo predictor can in principle cover a distinctly wider range than the Stambaugh Correlation, as the following proposition shows:

Proposition 3 *If x_t is the true predictor in the predictive regression (1), with predictive R-squared R_x^2 , and x_t^f is the fundamental pseudo predictor, which, from (9) can be constructed from the history of returns, then*

$$\text{corr}(x_t, x_t^f)^2 = \frac{R_f^2}{R_x^2} \geq \frac{R_f^2}{R_n^2} = \frac{1}{\theta^2} \left(\frac{\theta - \lambda}{\theta^{-1} - \lambda} \right)^2$$

where $R_f^2(\lambda, \theta)$ is the predictive R-squared of the fundamental ARMA representation, and equivalently of the fundamental pseudo predictor, and $R_n^2(\lambda, \theta)$ is R-squared of the non-fundamental representation

Proof. See Appendix F. ■

By inspection of the relationship in Proposition 3, it is evident that in the limiting case of white noise returns the correlation is precisely zero, since this implies $R_f^2 = 0$. Indeed it is of the essence of a white noise process that its own history is entirely uninformative about its own future values, and hence it must be uninformative about any predictor of its future values.

²⁶To complicate matters further, in Robertson & Wright, 2009b we show that it also an endemic feature of predictors that are actually redundant once we correctly condition on the history of returns, thus making it hard to distinguish between true and redundant predictors, especially if returns exhibit significant variance compression.

For near-white noise processes the correlation is non-zero, and the proposition shows that the better the true predictor predicts, the less similar it will be to the fundamental pseudo predictor. But the upper bound on R_x^2 given in Proposition 1 implies a *lower* bound on the correlation in Proposition 3: hence the narrower is the range of possible values of R_x^2 , the more similar the true predictor must be to the fundamental pseudo predictor. The lower bound in Proposition 3 is determined by the relative predictive power of the fundamental vs non-fundamental representations.

Figure 6 (below Figure 5) illustrates, for the three near-white noise processes already illustrated in Figure 5. Since all three have the same value of R_f^2 , the univariate R-squared, the relationship between $\text{corr}(x_t, x_t^f)$ and R_x^2 given in Proposition 3 is identical for all three processes. If the true predictor predicts barely any better than the univariate representation, it will very closely resemble it, but the better it predicts the weaker this resemblance will be. The only impact of greater variance compression (a lower value of V) will be that, since this reduces the upper bound for R_x^2 , it must *increase* the lower bound for $\text{corr}(x_t, x_t^f)$.²⁷

However, Figure 6 illustrates that even when there is very significant variance compression (as θ approaches unity) there is still scope for the true predictor to look quite *dissimilar* to the fundamental pseudo predictor. This suggests a simple pre-test when looking for predictor variables for stock returns: we should seek those that do *not* simply look like the history of returns.

5 The predictive space for real US stock returns: some empirical illustrations

5.1 A special case: white noise

We first consider the simplest possible case, in which returns are assumed to be entirely unpredictable from their own past. This is worth considering not just as a benchmark for comparison, but also because it is an implication (whether implicit or explicit) of a range of revisionist inves-

²⁷Figure 5 is placed below Figure 4 to illustrate that, for each process in Figure 5, the lower bound corresponds to the point in Figure 4 where the Stambaugh correlation for that process hits unity.

tigations of return predictability. Some of these have concluded that there is simply no return predictability at all of any kind (eg, Goyal & Welch, 2003); others (Kim et al, 1991, Pastor & Stambaugh, 2008) have concluded that the true variance ratio does not differ significantly from unity at any horizon, which must imply directly that there is no univariate predictability. Even defenders of return predictability such as Campbell & Viceira (1999) have acknowledged the logical possibility that there may be no univariate predictability.

Of course if returns are white noise, we have no way of inferring anything directly from the history of returns about the values of the ARMA parameters, except that they must be equal. But this does not mean we can infer nothing about the predictive space: it simply means that its properties depend on a single parameter, λ , the persistence of the true predictor, since the white noise assumption means that the predictive space must always satisfy $\theta(\lambda, \rho, R_x^2) = \lambda$. We noted in Section 2 that we have *a priori* grounds for assuming that predictors are likely to be persistent. In Table 2 we show that the maximum possible predictive R-squared, as given by Proposition 1, declines as λ increases²⁸. For reasonably persistent predictors the scope for return predictability, from *any* possible predictor of a white noise return process, is therefore quite limited.

As noted in our discussion of Proposition 2, the white noise case means that in principle the Stambaugh Correlation, ρ , can live anywhere in $(0, 1)$; but a *useful* predictor must have a positive Stambaugh Correlation, and the better it predicts the higher ρ must be (since the limiting case of the best possible predictor is the non-fundamental pseudo predictor defined in Section 3.2.2, with $\rho = 1$). Furthermore, since a higher value of λ brings down the upper bound at which ρ reaches unity, for any given value of R_x^2 , ρ is also increasing in λ .

The bottom row of Table 2 illustrates, by showing the required value of ρ consistent with $R_x^2 = 0.05$ - a figure not out of line with those found in the return predictability literature.

Table 2. The predictive space if stock returns are white noise

| | | | | | | | | |
|---|-----|------|------|------|------|------|------|------|
| λ | 0.0 | 0.25 | 0.5 | 0.6 | 0.7 | 0.8 | 0.8 | 0.95 |
| $\max(R_x^2) = R_n^2(\lambda, \lambda)$ | 1 | 0.94 | 0.75 | 0.64 | 0.51 | 0.36 | 0.19 | 0.10 |
| $\rho R_x^2 = 0.05$ | 0 | 0.06 | 0.13 | 0.17 | 0.22 | 0.31 | 0.47 | 0.70 |

²⁸Given the white noise assumption it is simply given by $\max(R_x^2) = R_n^2(\lambda, \lambda) = 1 - \lambda^2$

The necessary link between ρ and R_x^2 in the case of white noise returns casts an interesting light on the predictability literature. As noted above, valuation ratios such as the price-dividends and price-earnings ratios have frequently been proposed as predictors of returns. In Robertson & Wright (2009b) we show that a range of such predictors all have AR(1) parameters in the neighbourhood of 0.9; but the estimated Stambaugh Correlations associated with these predictors are *also* around this value, or in some cases, even closer to unity, and thus, as Table 2 shows, are very much higher than would be consistent with returns being white noise. An immediate conclusion that follows is that it would not be possible to claim simultaneously that any one of these predictors is the true predictor, *and* that returns are white noise. We shall see in the next section that such high values of ρ *are* more consistent with a predictor of a return process with a declining variance ratio, but in that case the univariate predictability that necessarily follows from this provides an alternative benchmark against which to compare such predictors. In Robertson & Wright (2009b) we conclude that none of these commonly used predictors can be distinguished in the data from the pseudo predictor consistent with this univariate predictability.

5.2 Estimating the predictive space for US stock returns, 1871-2008

If we move away from the white noise case, by allowing for the possibility that returns may be near-white noise with variance compression, it follows straightforwardly that the predictive space must contract. Variance compression requires $\theta > \lambda$. This raises R_f^2 , the fit of the fundamental ARMA representation, above unity, but at the same time *decreases* the maximum possible predictive R-squared, given by the non-fundamental R-squared, R_n^2 , thus contracting the space that R_x^2 can feasibly inhabit. At the same time, from Propositions 2 and 3, increasing variance compression raises towards unity the lower bounds on both the Stambaugh Correlation and the correlation between the predictor and the fundamental pseudo predictor. Thus on the basis of a priori reasoning alone we know that greater is the degree of variance compression, the more the predictive space must contract.

In estimating the limits to the predictive space consistent with the observed history of

returns we should note at the outset that, given the near-white noise properties of returns, no method of estimation can be expected to yield well-determined results. Nor do we wish to pin ourselves down to any assumption that the univariate representation has been stable, and of the restrictive ARMA(1,1) form, over the entire sample of returns.²⁹ Our estimates in this section are thus largely illustrative.

A starting point is simply to estimate the ARMA representation. There are obvious caveats: the near-white noise property means that the AR and MA components are very close to cancellation, and thus, as is well known, both λ and θ are likely to be poorly estimated, and subject to significant small-sample (essentially Stambaugh, 1999) bias. There is however an important cross-check on our results, in the spirit of Cochrane, 1988. We showed in Section 3.3 that in the ARMA(1,1) there is a direct correspondence between the sign of $\theta - \lambda$ and the slope of the variance ratio. It is also straightforward to show³⁰ that, if $\theta > \lambda$, the *rate* at which the variance ratio slopes downwards is determined solely by the magnitude of $V(\lambda, \theta)$ (the limiting variance ratio) and λ . In principle direct measurement of the variance ratio could, for some processes, yield very different answers from that implied by ARMA estimates;³¹ but in both the long annual sample 1871-2008 and (with caveats) the shorter postwar sample 1945-2008, the results are reassuringly similar.

Figure 7 illustrates. We estimate ARMA(1,1) representations of returns in both samples. In terms of the expectations derived from our analysis thus far the point estimates are certainly in the right ballpark: for the full sample we have $\hat{\lambda} = 0.860$ and $\hat{\theta} = 0.977$, and in the postwar sample we have $\hat{\lambda} = 0.89$ $\hat{\theta} = 0.95$, thus in both samples the point estimates are consistent with variance compression, but they are somewhat closer together in the postwar sample, and hence somewhat closer to white noise. Figure 7 shows that if we treat the ARMA estimates as equal to their true population values, the results are not in conflict with the evidence from


²⁹It is quite possible that there may have been shifts in the ARMA parameters as well as both the volatility and unconditional mean of real returns. However, there is a non-trivial caveat: the *extent* of any shifts in the mean return cannot have been too large. If such shifts are treated as stochastic, in, for example, a Markov Switching Model, Hamilton (1989) showed that that the reduced form process has an ARMA representation. For sufficiently strong shifts it is possible to show that the declining variance ratio observed in the data would be ruled out.

³⁰See Appendix C.

³¹See, for example, the comparison between the very different implications of the variance ratio and ARMA representations of GDP growth in Cochrane, 1988.

direct measurement of the variance ratio.

**Figure 7. The Variance Ratio:
Data vs Implications of ARMA Estimates³²**



In the full sample this consistency is particularly marked. The implied “true” horizon variance ratio matches the sample variance ratio well, particularly at longer investor horizons; and even when the two profiles differ somewhat at shorter horizons, the deviation is well within the range of sampling variation.³³

In the post-war sample, while the ARMA estimates are quite similar to those estimated over the full sample, they are less consistent with direct measurement of the variance ratio. But the differences are not in general statistically significant. Given the short sample, if the estimated ARMA parameters were truly generating the returns data, the range of sampling variation of the variance ratio would be quite wide, especially at short horizons, hence the lack of any decline for horizons up to around 15 years (as noted by Kim et al, 1991) would not of itself be particularly significant. Indeed the only statistically significant contrast between the

³²Figure 7 shows the variance ratio in the data, as in Figure 2, but the 5% and 95% bounds and mean estimates are simulated in 10,000 replications using the estimated ARMA model as the data generating process. The two panels also show the true value of the variance ratio, as given by (34) in Appendix C, on the same assumptions.

³³Note that if this is the true data generating process the extent of sampling variation in the directly measured variance ratio is very much lower than in the white noise case, additionally there is essentially no small sample bias.

two approaches is at very long horizons, when the observed variance ratio actually breaches the lower 5% bounds consistent with the ARMA estimates being the true model. However, given the range of uncertainty in both approaches, it is fairly obvious that it would take only a very limited amount of data mining to find an ARMA representation that was consistent both with the direct ARMA estimates and the evidence of the variance ratio, over both samples. Any such representation would have a high value of λ , and $\theta > \lambda$.

**Table 3 Point Estimates of Limits on the Predictive Space for US Stock Returns
Implied by ARMA Estimates³⁴**

| | (1) | (2) | (3) | (4) | (5) | (6) |
|-----------|-----------------|----------------|---------------|---------------|---------------|---------------------------------|
| Sample | $\hat{\lambda}$ | $\hat{\theta}$ | $\min(R_x^2)$ | $\max(R_x^2)$ | ρ_{\min} | $\min[\text{corr}(x_t, x_t^f)]$ |
| 1871-2008 | 0.860 | 0.977 | 0.052 | 0.095 | 0.986 | 0.739 |
| 1945-2008 | 0.891 | 0.955 | 0.020 | 0.106 | 0.908 | 0.429 |

Given the mutual consistency of the two approaches (particularly in the long sample) we have no obvious reason, in terms of the variance ratio evidence at least, to object to the ARMA estimates. In Table 3 we therefore take these estimates at face value, and use them to calculate the implied constraints on the predictive space, using estimates from both long and short samples.

The implied value of $\min(R_x^2) = R_f^2$ the univariate R-squared, which, from Proposition 1, provides the lower bound for the predictive R-squared of the true predictor is around 5% in the long sample. This is reasonably consistent with the sample estimate (if anything, given the known impact of Stambaugh Bias, we might expect the sample value to be rather higher). In the postwar sample, as noted above, returns appear closer to white noise, hence the implied true R_f^2 is distinctly closer to zero.³⁵

³⁴Columns (1) and (2) show the estimated autoregressive (λ) and moving average (θ) parameters in estimated ARMA(1,1) representations of returns over the given samples. Columns (3) and (4) give the implied upper and lower bounds for the predictive R-squared from Proposition 1, given by (column 3) $\min(R_x^2) = R_f^2(\hat{\lambda}, \hat{\theta})$ and (column 4) $\max(R_x^2) = R_n^2(\hat{\lambda}, \hat{\theta})$. Column (5) gives the implied lower bound, $\rho_{\min}(\hat{\lambda}, \hat{\theta})$ for the Stambaugh Correlation from Proposition 2. Column (6) gives the lower bound for the correlation between the true predictor and the pseudo predictor, as given by Proposition 3, as $\sqrt{R_f^2(\hat{\lambda}, \hat{\theta}) / R_n^2(\hat{\lambda}, \hat{\theta})}$.

³⁵Note that in our theoretical analysis we focussed on the true R^2 , which is a function of the true values of

For the upper bound, $\max(R_x^2) = R_n^2$, the notional R-squared of the non-fundamental ARMA representation, we have, of course, no cross-check from the data, but we can estimate it directly from the estimated values of λ and θ if we treat them as the true parameters. This calculation implies that, in both samples, the best possible predictor of stock returns would have an R-squared of around 10%: thus in terms of predictive R-squared the predictive space is quite narrow. The implied space for the Stambaugh Correlation is even more tightly constrained: the point estimate of ρ_{\min} , as defined in Proposition 2, is very close indeed to unity, particularly using full sample estimates.

In the final column of the table we calculate the implied lower bound for the correlation between the true predictor and the fundamental pseudo predictor. It is noticeable that, despite the apparently very limited predictive space for R_x^2 and ρ , the true predictor can still in principle look reasonably different from the pseudo predictor - particularly so if we use the postwar ARMA estimates.³⁶ Nonetheless the clear implication of Table 3 is that point estimates consistent with the data suggest only very little space for any predictor to out-predict the univariate representation.

Of course, given the known problems in estimating ARMA representations with near-cancellation of AR and MA roots, the figures in Table 3 cannot be treated as anything other than illustrative. We certainly would not wish to state categorically that the true predictive space must be as narrow as the ARMA estimates suggest. Given sampling variation, the history of returns is in principle consistent with a range of true data generating processes, some of which have a distinctly less constrained predictive space. However, if we wish to argue that the predictive space *is* less constrained, we show in the next section that this has important implications for another aspect of return predictability on which we have not yet touched: namely long-horizon predictability.

λ and θ . In any given finite sample if we use the formula for the population value to calculate $R_f^2(\hat{\lambda}, \hat{\theta})$ the result of this calculation need not be the same as the sample R-squared calculated from the ARMA estimation; although in practice simulation evidence shows that the two figures are usually quite close.

³⁶This reflects the fact that, while the difference between R_f^2 and R_n^2 is small, it is the *ratio* that determines the lower bound for the correlation, and in the postwar sample in particular this ratio is quite high, despite the low value of R_n^2 itself.

6 The predictive space and long-horizon return predictability

We have stressed already that the estimates in Table 3 are only illustrative. In the Appendix we show that given the range of sampling variation of ARMA estimates we might quite easily derive point estimates of a similar order of magnitude to those in Table 3 for a quite wide range of white and near-white noise processes. Indeed if we focus purely on statistical criteria, the only clear-cut results are:

- First, and fairly obviously, the true process cannot be very far from white noise. In terms of Figure 4, the true values of θ and λ must lie within the quite narrow range given by the R-squared contour lines. Hence we would reject any data generating process for returns for which θ was very far from λ .
- Additionally we have even stronger grounds to reject near-white noise processes with variance *expansion* rather than compression (ie, with $\theta < \lambda$). Even if the true variance ratio has only a modest upward slope, the probability of observing the low values observed in the data rapidly becomes vanishingly small.
- On the other hand we know that the rejection of the strict white noise case is at best only marginal. Hence the data also do not reject values of θ and λ for which the variance ratio only slopes down very modestly (ie, for which V is less than, but quite close to unity)

While these considerations rule out quite a wide range of (λ, θ) combinations, it is evident that the data *do* admit representations that lie roughly between the 45° line and the upper R-squared contour in Figure 4. Since this includes representations in which λ and θ are both close to zero, this means that predictive space could be very much less constrained than the point estimates in Table 3 suggest. For true processes sufficiently close to white noise, and with sufficiently low λ , we showed in Section 5.1, Table 2, that the upper bound for R_x^2 could in principle be close to unity, and the lower bound for ρ could be close to (though not below) zero.

However, while this is a logical possibility, if we *did* wish to argue that the predictive space were significantly wider, we could only do so by simultaneously assuming that another often-assumed characteristic of stock returns is absent, namely long-horizon predictability.

Any predictive regression system has an associated profile for the horizon R-squared, given by $R_x^2(h)$, which is the R-squared in predicting the average return over h periods from period t , $r_{t,t+h} \equiv h^{-1} \sum_{i=1}^h r_{t+h}$. A convenient way to summarise the characteristics of this horizon profile is to define the horizon, h^* at which the horizon R-squared is at its maximum value. In Appendix G we show that the system in (1) and (2) has the convenient property that the horizon R-squared profile for the true predictor is simply a scaling of the univariate horizon profile. It follows that we have

$$h^* \equiv \arg \max (R_x^2(h)) = \arg \max (R_f^2(h)) = h^*(\lambda, \theta) \quad (20)$$

so that, while the true predictor could in principle have a very different value of $R_x^2(h^*)$ from the univariate representation, the horizon at which this maximum occurs is identical to that for the equivalent univariate R-squared profile, $R_f^2(h)$.³⁷

Table 4 illustrates the link between horizon predictability and different (λ, θ) pairs in the true process for returns. The shaded area shows values for which the one-period ahead univariate R-squared is less than 5%, hence any data-consistent representation of returns must lie roughly in this area.

The table shows that for quite a wide range of (λ, θ) pairs close to zero h^* is unity. For such combinations the horizon R-squared profile declines monotonically, hence there are no horizon effects at all. Only as we move north-westwards, for relatively high values of both parameters, do horizon effects become more significant. Furthermore, for given λ , the higher is θ (hence the stronger is variance compression, the stronger the horizon effects) the But we have seen already that is precisely for these parameter values that the predictive space contracts. Hence there is a near equivalence between strong horizon effects and a tightly constrained predictive space.

³⁷The expression in (20) holds for any value of $\theta \neq \lambda$, hence for any return process that is arbitrarily close to white noise. For the case $\theta = \lambda$ we can however still derive the same property for h^* , by using the horizon profile of the non-fundamental pseudo predictor, $R_n^2(h)$.

Or, putting it another way, if there *is* strong long-horizon predictability, it must be close to being a univariate phenomenon.

Table 4. The optimal horizon for the horizon R-squared of the true predictor of ARMA(1,1) returns.³⁸

| | | AR Parameter, λ | | | | | | | |
|------------------------|------|-------------------------|-----|-----|-----|-----|-----|------|------|
| | | 0 | 0.2 | 0.4 | 0.6 | 0.8 | 0.9 | 0.95 | 0.98 |
| MA Parameter, θ | 0.98 | 1 | 5 | 8 | 13 | 22 | 33 | 46 | 62 |
| | 0.95 | 1 | 4 | 6 | 9 | 14 | 20 | 24 | 29 |
| | 0.9 | 1 | 3 | 5 | 7 | 9 | 12 | 14 | 15 |
| | 0.8 | 1 | 2 | 3 | 4 | 6 | 7 | 7 | 7 |
| | 0.6 | 1 | 2 | 2 | 2 | 3 | 3 | 3 | 3 |
| | 0.4 | 1 | 1 | 1 | 2 | 2 | 2 | 2 | 2 |
| | 0.2 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 0 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

7 Conclusions

This paper shows that the univariate properties of stock returns can be used to infer restrictions on the nature of any true predictor variable for stock returns, because these univariate properties depend in specific ways on the parameters that characterise the predictive system, which we call the predictive space. The argument does not hinge particularly on whether or not we are able to estimate a univariate ARMA model precisely, because merely knowing that returns are near serially uncorrelated and have a downward sloping variance ratio provides sufficient information to restrict strongly the predictive space.

Our results have three strong implications for the return predictability literature. First, if predictor variables are persistent then it may be that the predictive space contracts to such an extent that no predictor variable will predict very much better than a pseudo-predictor, that is itself simply a weighted average of past returns. Second, the converse also applies, that is if we seek predictor variables that have meaningful predictive ability, they will have to be dissimilar

³⁸The table shows the value of h^* , as defined in equation (20) for different values of θ and λ in the ARMA(1,1) representation of returns in (3). The figures for h^* shown in the table are calculated numerically using the general formula for the horizon R-squared profile in (41) in Appendix G. The shaded area shows (λ, θ) pairs with $R_T^2 \leq 0.05$, bolded figures are white noise.

to weighted past returns and will likely have lower persistence than most existing predictor variables. Third, strong horizon effects imply a tightly constrained predictive space, leaving little scope for the true predictor to outperform the univariate representation.

Appendix

A Deriving the ARMA(1,1) structure.

Start from

$$r_t = -\beta x_{t-1} + u_t$$

$$x_t = \lambda x_{t-1} + v_t$$

where $\begin{pmatrix} u_t \\ v_t \end{pmatrix}$ are jointly serially uncorrelated mean zero with covariance matrix $\begin{pmatrix} \sigma_u^2 & \sigma_{uv} \\ \sigma_{uv} & \sigma_v^2 \end{pmatrix}$.

Solving gives us the ARMA reduced form for r_t

$$r_t = \lambda r_{t-1} + \varepsilon_t - \theta \varepsilon_{t-1}$$

where

$$\varepsilon_t - \theta \varepsilon_{t-1} = u_t - \lambda u_{t-1} - \beta v_{t-1}$$

Where θ must satisfy the moment condition

$$\begin{aligned} \frac{-\theta}{1 + \theta^2} &= \frac{\text{cov}(-\beta v_{t-1} + (1 - \lambda L)u_t, -\beta v_{t-2} + (1 - \lambda L)u_{t-1})}{\text{var}(-\beta v_{t-1} + (1 - \lambda L)u_t)} \\ &= \frac{-(\lambda \sigma_u^2 + \beta \sigma_{vu})}{\beta^2 \sigma_v^2 + \sigma_u^2 (1 + \lambda^2) + 2\lambda \beta \sigma_{vu}} = \frac{-(\lambda + \beta \rho s)}{1 + \lambda^2 + \beta^2 s^2 + 2\lambda \rho \beta s} \end{aligned}$$

where $\rho = \sigma_{vu}/(\sigma_u \sigma_v)$; $s = \sigma_v/\sigma_u$.

Note also that if R_x^2 is the R-squared from the predictive regression then

$$R_x^2 = \frac{\beta^2 \sigma_x^2}{\sigma_r^2} = \frac{\beta^2 s^2}{\beta^2 s^2 + 1 - \lambda^2}$$

implying

$$\beta^2 s^2 = F^2 \text{ where } F(R_x^2, \lambda) = \sqrt{(1 - \lambda^2) \frac{R_x^2}{1 - R_x^2}} \quad (21)$$

hence the moment condition defining θ can be written as

$$\frac{\theta}{1 + \theta^2} = \kappa(\lambda, \rho, R_x^2) \quad (22)$$

$$\text{where } \kappa(\lambda, \rho, R_x^2) = \frac{\lambda + \rho F(R_x^2, \lambda)}{1 + \lambda^2 + F(R_x^2, \lambda)^2 + 2\lambda\rho F(R_x^2, \lambda)} \quad (23)$$

which solves to give the MA parameter in the fundamental representation³⁹

$$\theta(\lambda, \rho, R_x^2) = \frac{1 - (1 - 4\kappa(\lambda, \rho, R_x^2)^2)^{\frac{1}{2}}}{2\kappa(\lambda, \rho, R_x^2)} \quad (24)$$

A real solution for $\theta \in (-1, 1)$ requires that $\kappa \in (-\frac{1}{2}, \frac{1}{2})$. To show this holds, note first that we have

$$\partial\kappa/\partial\rho = \frac{F(1 + F^2 - \lambda^2)}{(1 + \lambda^2 + F^2 + 2\lambda\rho F)^2} > 0 \quad (25)$$

for $\lambda \in (0, 1)$, $R_x^2 \in (0, 1)$. Thus, since $\rho \in (-1, 1)$, we know that

$$\begin{aligned} \kappa(\lambda, \rho, R_x^2) &\in (\kappa(\lambda, -1, R_x^2), \kappa(\lambda, 1, R_x^2)) \\ &\in (g(\lambda - F), g(\lambda + F)) \end{aligned}$$

where

$$g(x) = \frac{x}{1 + x^2} \in \left(-\frac{1}{2}, \frac{1}{2}\right) \quad (26)$$

hence we do indeed have $\kappa \in (-\frac{1}{2}, \frac{1}{2})$. Given this, we know that

$$\frac{\partial\theta}{\partial\kappa} = \frac{1}{2} \left(\frac{1 - \sqrt{1 - 4\kappa^2}}{\kappa^2 \sqrt{1 - 4\kappa^2}} \right) \geq 0 \quad (27)$$

which in turn gives us

$$\frac{\partial\theta}{\partial\rho} = \frac{\partial\theta}{\partial\kappa} \frac{\partial\kappa}{\partial\rho} \geq 0 \quad (28)$$

which we shall exploit later.

To derive the representation in terms of the pseudo predictor in (5) and (6), write the

³⁹The other solution to (23) gives the nonfundamental representation.

ARMA as

$$r_t = \left(\frac{1 - \theta L}{1 - \lambda L} \right) \varepsilon_t = \varepsilon_t + \frac{(\lambda L - \theta L)\varepsilon_t}{1 - \lambda L} = \varepsilon_t + (\lambda - \theta) \frac{\varepsilon_{t-1}}{1 - \lambda L}$$

if we then define the pseudo predictor as

$$x_t^f = \text{sign}(\theta - \lambda) \frac{\varepsilon_t}{1 - \lambda L}$$

and substitute we can write the system as in (5) and (6), with $\beta_f = |\theta - \lambda|$ and $\rho = \text{sign}(\theta - \lambda)$, which is nested within the general system (1) and (2), preserving the useful convention that β_x is always negative

B The Fundamental ARMA R-squared

We have

$$R_f^2 = 1 - \frac{\sigma_\varepsilon^2}{\sigma_r^2}$$

We can use the Yule-Walker equations to derive

$$\sigma_r^2 = \left(\frac{1 - \lambda^2 + (\theta - \lambda)^2}{1 - \lambda^2} \right) \sigma_\varepsilon^2$$

hence

$$R_f^2 = 1 - \frac{\sigma_\varepsilon^2}{\sigma_r^2} = \frac{(\theta - \lambda)^2}{1 - \lambda^2 + (\theta - \lambda)^2} \quad (29)$$

C The Variance Ratio and ARMA Parameters

The standard definition of the variance ratio given in Cochrane (1988) is

$$VR(h) = 1 + 2 \sum_{j=1}^{h-1} \left(\frac{h-j}{h} \right) \text{corr}(j) \quad h = 1, 2, \dots \quad (30)$$

where $corr(h) = corr(r_t, r_{t-h})$. In the ARMA(1,1) we have, using the Yule-Walker equations,

$$corr(1) = \frac{cov(r_t, r_{t-1})}{var(r_t)} = - \left(\frac{(\theta - \lambda)(1 - \theta\lambda)}{1 - \lambda^2 + (\theta - \lambda)^2} \right) \quad (31)$$

$$corr(j) = corr(1)\lambda^{j-1}; \quad j > 1 \quad (32)$$

hence substituting and evaluating the sum we have

$$VR(h) = 1 + 2 \cdot \frac{corr(1)}{1 - \lambda} \left(1 - \frac{1 - \lambda^h}{h(1 - \lambda)} \right) \quad (33)$$

and note that $\frac{1 - \lambda^h}{h}$ is decreasing in h for $h > 1$ so $VR(h)$ decreases or increases monotonically from $VR(1) = 1$.

As $h \rightarrow \infty$ we have the limiting variance ratio

$$\lim_{h \rightarrow \infty} VR(h) = 1 + 2 \cdot \frac{corr(1)}{1 - \lambda} = 1 - \frac{2}{1 - \lambda} \left(\frac{(\theta - \lambda)(1 - \theta\lambda)}{1 - \lambda^2 + (\theta - \lambda)^2} \right)$$

noting that

$$1 - R_f^2 = \frac{1 - \lambda^2}{1 - \lambda^2 + (\theta - \lambda)^2} = \frac{(1 - \lambda)(1 + \lambda)}{1 - \lambda^2 + (\theta - \lambda)^2}$$

and writing

$$\begin{aligned} V &= \lim_{h \rightarrow \infty} VR(h) = \frac{1}{1 - \lambda} \left(\frac{(1 - \lambda)(1 - \lambda^2 + (\theta - \lambda)^2) - 2(\theta - \lambda)(1 - \theta\lambda)}{1 - \lambda^2 + (\theta - \lambda)^2} \right) \\ &= \frac{(1 - R_f^2)((1 - \lambda)(1 - \lambda^2 + (\theta - \lambda)^2) - 2(\theta - \lambda)(1 - \theta\lambda))}{(1 - \lambda)^2(1 + \lambda)} \end{aligned}$$

which simplifies to

$$V = (1 - R_f^2) \left(\frac{1 - \theta}{1 - \lambda} \right)^2$$

Now by inspection

$$\theta > \lambda \Rightarrow V < 1$$

Conversely

$$\begin{aligned}
V < 1 &\Rightarrow \\
\left(\frac{1 - \lambda^2}{1 - \lambda^2 + (\theta - \lambda)^2} \right) (1 - \theta)^2 &< (1 - \lambda)^2 \\
0 &< \lambda^2 - \lambda - \lambda\theta^2 + \lambda^2\theta^2 - \theta\lambda + \lambda^2\theta - \lambda^3\theta + \theta \\
0 &< (1 - \lambda^3)\theta + (1 + \theta + \theta^2)(\lambda^2 - \lambda) \\
(1 + \theta + \theta^2)\lambda(1 - \lambda) &< (1 - \lambda^3)\theta \\
\frac{\lambda(1 - \lambda)}{1 - \lambda^3} &< \frac{\theta}{1 + \theta + \theta^2} = \frac{\theta(1 - \theta)}{1 - \theta^3} \\
f(\lambda) &< f(\theta)
\end{aligned}$$

with $f(x) = \frac{x(1-x)}{1-x^3}$. Now $f(\cdot)$ is a monotone increasing function since

$$f'(x) = \frac{1 - x^2}{(1 + x + x^2)^2} > 0 \text{ for } |x| < 1$$

hence

$$V < 1 \Leftrightarrow \theta > \lambda$$

and this plus monotonicity of $VR(h)$ gives the result in (17).

Finally note that we have

$$VR(h) = V + (1 - V)\sqrt{\frac{H(h, \lambda)}{h}} \tag{34}$$

where

$$H(h, \lambda) = \frac{1}{h} \left(\frac{1 - \lambda^h}{1 - \lambda} \right)^2$$

D Proof of Proposition 1

We wish to establish the inequality

$$R_f^2(\lambda, \theta) \leq R_x^2 < R_n^2(\lambda, \theta)$$

D.1 Relation of R_x^2 to R_f^2

The first inequality is straightforward. Using the derivation of the ARMA(1,1) representation in Appendix A we have

$$\begin{aligned}\varepsilon_t &= \frac{1}{1 - \theta L} [-\beta v_{t-1} + (1 - \lambda L)u_t] \\ &= u_t - \lambda u_{t-1} - \beta v_{t-1} + \theta \varepsilon_{t-1} \\ &= u_t + \psi_{t-1}\end{aligned}$$

hence

$$\text{var}(\varepsilon_t) = \text{var}(u_t) + \text{var}(\psi_t) > \text{var}(u_t)$$

since $\text{cov}(u_t, \psi_{t-1}) = 0$. Hence for $1 < \rho < 1$ we always have

$$\begin{aligned}1 - \frac{\sigma_\varepsilon^2}{\sigma_r^2} &< 1 - \frac{\sigma_u^2}{\sigma_r^2} \\ R_f^2 &< R_x^2\end{aligned}$$

For the fundamental pseudo predictor case, if $\theta > \lambda$ we have $u_t = \varepsilon_t = v_t$ (hence $\rho = 1$) so $\psi_{t-1} = (\theta - \lambda - \beta) \varepsilon_{t-1} = 0$ and if $\theta < \lambda$ we have $u_t = \varepsilon_t = -v_t$ (hence $\rho = -1$), so $\psi_{t-1} = (\theta - \lambda + \beta) \varepsilon_{t-1} = 0$. Hence for the limiting case of the fundamental pseudo predictor we always have $\sigma_u^2 = \sigma_\varepsilon^2 \Rightarrow R_f^2 = R_x^2$ so for the general case we have

$$R_f^2 \leq R_x^2$$

D.2 Relation of R_x^2 to R_n^2

We have the non-fundamental representation

$$r_t = \left(\frac{1 - \theta^{-1}L}{1 - \lambda L} \right) \eta_t \tag{35}$$

where η_t is the non-fundamental innovation, and we know (Hamilton, 1994, pp 66-67)

$$\sigma_\eta^2 = \theta^2 \sigma_\varepsilon^2$$

hence

$$R_n^2 = R_f^2 + (1 - \theta^2) \frac{\sigma_\varepsilon^2}{\sigma_r^2} = R_f^2 + (1 - \theta^2) (1 - R_f^2)$$

which, after substituting from (29) gives

$$R_n^2 = \frac{(1 - \theta\lambda)^2}{1 - \lambda^2 + (\theta - \lambda)^2} \quad (36)$$

which can also be derived directly from (29) by substituting θ^{-1} for θ .

Note that for $\theta = 0$ the non-fundamental representation (10) is undefined. As $\theta \rightarrow 0$ we have $\sigma_\eta^2 = \theta\sigma_\varepsilon^2 \rightarrow 0$, but, expanding (11),

$$E_t r_{t+1} | x_t^n = -(\theta^{-1} - \lambda) [-\theta r_{t+1} - \theta^2 r_{t+2} + \dots]$$

so we have $\lim_{\theta \rightarrow 0} E_t r_{t+1} | x_t^n = r_{t+1}$, hence $\lim_{\theta \rightarrow 0} R_n^2 = 1$.

We wish to establish the inequality

$$G(\lambda, \rho, R_x^2) \equiv R_n^2(\lambda, \theta(\lambda, \rho, R_x^2)) - R_x^2 \geq 0$$

While G depends in principle on the triplet (λ, ρ, R_x^2) we shall analyse its properties for a given (λ, R_x^2) pair; we shall show that the result hold for any (λ, R_x^2) within their allowable ranges. Note that for this proof we do not require λ to be positive.

Thus we can write $G = G(\theta(\rho))$. From (28) we also have $\partial\theta/\partial\rho > 0$, hence we can write

$$G = G(\theta); \quad \theta \in [\theta_{\min}, \theta_{\max}]$$

where

$$\theta_{\min} = \theta(\lambda, -1, R_x^2); \quad \theta_{\max} = \theta(\lambda, 1, R_x^2)$$

and we have

$$\begin{aligned} G'(\theta) &= \frac{\partial R_n^2}{\partial \theta} = -2\theta \left(\frac{(1-\lambda^2)(1-\theta\lambda)}{1-\lambda^2+(\theta-\lambda)^2} \right) \\ &\Rightarrow \text{sign}(G'(\theta)) = -\text{sign}(\theta) \end{aligned} \quad (37)$$

There are three cases:

Case 1: $\theta_{\min} > 0$; For this case, we have $G(\theta_{\max}) = 0$, since at this point x is the non-fundamental pseudo predictor with $R_x^2 = R_n^2$, $\rho = \rho_n = 1$. From (37) we also have $G' < 0$ hence $G \geq 0$.

Case 2: $\theta_{\max} < 0$; For this case, we have $G(\theta_{\min}) = 0$, since at this point x is again the non-fundamental pseudo predictor with $R_x^2 = R_n^2$, but with $\rho = \rho_n = -1$. From (37) we have $G' > 0$ hence $G \geq 0$.

Case 3: $\theta_{\min} < 0 < \theta_{\max}$; For this case, we have $G(\theta_{\min}) = G(\theta_{\max}) = 0$, $G'(\theta_{\min}) > 0$; $G'(\theta_{\max}) < 0$, and, from (37) G has a single turning point at $\theta = 0$, hence again we have $G \geq 0$.

Since we have shown that these results hold for any θ and λ , by implication they also hold for any λ and R_x^2 .

We have thus established the right-hand inequality in (19) for all three cases thus completing the proof. ■

E Proof of Proposition 2

Any given values of θ and λ must imply a condition on κ (as defined in (23)) of the form

$$\kappa(\lambda, \rho, R_x^2) = \frac{\theta}{1 + \theta^2}$$

For given values of θ and λ this can be taken to imply a restriction on ρ , the correlation between the two underlying innovations, which solves to give

$$\rho(\theta, \lambda, R_x^2) = \frac{(\theta - \lambda)(1 - \theta\lambda) + F(\lambda, R_x^2)^2 \theta}{(1 - \lambda^2 + (\theta - \lambda)^2) F(\lambda, R_x^2)}; \quad \rho \in (-1, 1) \quad (38)$$

where $F(\lambda, R_x^2)$, as defined in (21). If the solved value for ρ lies outside this range, the triplet (θ, λ, R_x^2) is not feasible (ie, taking θ and λ as given, R_x^2 does not satisfy the inequality condition (19)).

The first order condition yields a unique stationary point:

$$\frac{\partial \widehat{\rho}(\theta, \lambda, R_x^2)}{\partial R_x^2} = 0 \Rightarrow R_x^2 = \frac{(\theta - \lambda)(1 - \theta\lambda)}{\theta - \lambda + \theta(1 - \theta\lambda)}$$

which after substituting into (38) yields a real solution if

$$(\theta - \lambda)\theta > 0$$

which is satisfied for $\theta > \lambda$, given $\lambda > 0$. The second-order condition confirms that for this range of parameter values this yields the minimum value

$$\rho_{\min} = \text{sign}(\theta - \lambda) \frac{2\sqrt{(\theta - \lambda)(1 - \theta\lambda)\theta}}{(1 - \lambda^2 + (\theta - \lambda)^2)} > 0. \blacksquare$$

F Proof of Proposition 3

Consider the regression

$$r_t = -\beta_x x_{t-1} - \beta_f x_{t-1}^f + u_t$$

where x_t^f is the fundamental pseudo predictor and x_t the predictor variable. Then given the true model (1) that generates the data, it must be that $\beta_f = 0$. Treating x_{t-1} as an omitted variable in the regression

$$r_t = -\beta_f x_{t-1}^f + w_t$$

then $w_t = -\beta_x x_t + u_t$.

Using the formula for omitted variable bias we can obtain population values via moment conditions as

$$-\beta_f = \frac{\text{Cov}(x^f, r)}{\text{Var}(x^f)} = \frac{\text{Cov}(x^f, -\beta_x x + u)}{\text{Var}(x^f)} = -\beta_x \frac{\text{Cov}(x^f, x)}{\sqrt{\text{Var}(x^f)\text{Var}(x)}} \frac{\sqrt{\text{Var}(x)}}{\sqrt{\text{Var}(x^f)}}$$

$$= -\beta_x \rho_{x^f x} \frac{\sigma_x}{\sigma_{x^f}}$$

where $\rho_{x^f x} = \text{corr}(x^f, x)$

$$\beta_f \frac{\sigma_{x^f}}{\sigma_r} = \rho_{x^f x} \beta_x \frac{\sigma_x}{\sigma_r}$$

Squaring both sides we get

$$\left(\beta_f \frac{\sigma_{x^f}}{\sigma_r} \right)^2 = \left(\beta_x \frac{\sigma_x}{\sigma_r} \right)^2 \rho_{x^f x}^2$$

or

$$R_f^2 = \rho_{x^f x}^2 R_x^2$$

Using this formula for the non-fundamental R^2 we get

$$R_f^2 = \rho_{x^f x^n}^2 R_{n^f}^2$$

where $\rho_{x^f x^n} = \text{Corr}(x_t^f, x_t^{n^f})$.

G The horizon R-squared

Using (1) and (2), we have

$$E_t r_{t+h} = -\beta_x E_t x_{t+h-1} = -\beta_x \lambda^{h-1} x_t$$

$$E_t \left(\frac{1}{h} \sum_{i=1}^h r_{t+i} \right) = -\frac{\beta_x}{h} \sum_{i=1}^h \lambda^{h-1} x_t = -\beta_h x_t$$

where

$$\beta_h = \frac{\beta_x}{h} \left(\frac{1 - \lambda^h}{1 - \lambda} \right) = \beta_x \sqrt{\frac{H(h, \lambda)}{h}} \quad (39)$$

so the horizon coefficient profile is just a scaling of the variance ratio profile in (34).

We also know, from the definition of the variance ratio,

$$\text{var} \left(\frac{1}{h} \sum_{i=1}^h r_{t+i} \right) = \frac{1}{h^2} h \cdot \sigma_r^2 \text{VR}(h) = \frac{\sigma_r^2}{h} \text{VR}(h) \quad (40)$$

hence

$$R_x^2(h) = \beta_x^2 \frac{\sigma_x^2}{\sigma_r^2} \frac{H(h; \lambda)}{VR(h)} = R_x^2 \frac{H(h; \lambda)}{VR(h)} \quad (41)$$

Note that $H(h; \lambda)$ and $VR(h)$ depend only on ARMA parameters and h . Since we could also derive an equivalent horizon profile for the fundamental pseudo predictor, replacing R_x^2 with R_f^2 , the horizon profile for the true predictor is a scaling of the univariate horizon profile and hence has the same turning point, $h^*(\lambda, \theta)$.

References

- Ang, A and G Bekaert (2007) “Stock Return Predictability: Is it There?” *Review of Financial Studies* 20(3):651-707
- Beveridge, S. and C R Nelson. (1981) “A new approach to decomposition of economic time series into permanent and transitory components with particular attention to measurement of the business cycle”. *Journal of Monetary Economics*, 7, pp151-174.
- Boudoukh J, Richardson M and R Whitelaw (2008) “The Myth of Long Horizon Predictability” *Review of Financial Studies* 21(4):1577-1605
- Campbell J Y and R J Shiller (1998) “Valuation Ratios and the Long-Run Stock Market Outlook” *Journal of Portfolio Management*, 24, pp11–26
- Campbell, J Y, and S Thompson (2008) “Predicting Excess Stock Returns Out of Sample: Can Anything Beat the Historical Average?”, *Review of Financial Studies* 2008 21(4):1509-1531
- Campbell, J Y., and L M. Viceira, (2002), *Strategic Asset Allocation: Portfolio Choice for Long-Term Investors* (Oxford University Press)
- Campbell, J. Y., and M. Yogo, (2006), “Efficient Tests of Stock Return Predictability,” *Journal of Financial Economics*, 81(1), 27–60.
- Cochrane, J.H (1988) “How big is the random walk in GDP?. ” *Journal of Political Economy*, v. 96, n. 5, p. 893-92

- Cochrane J H (1994) "Permanent and Transitory Components of GNP and Stock Prices " *The Quarterly Journal of Economics* Vol. 109, No. 1, pp. 241-265
- Cochrane J H (2008) "The Dog that Did Not Bark: A Defense of Return Predictability" *Review of Financial Studies* 2008, 1533-1575.
- Cowles, A C (1938) *Common-Stock Indices* Principia Press, Bloomington Indiana
- Fama E F and K R French (2002) "The Equity Premium", *Journal of Finance* 62, pp637-659.
- Fernandez-Villaverde, J, Rubio-Ramirez, F, Sargent, T and Watson, M (2007) "ABCs (and Ds) of Understanding VARs", *American Economic Review*, pp 1021-1026
- Ferson W E Simin T and S Sarkissian (2003) "Spurious regressions in Financial Economics?", *Journal of Finance* 58, 1393-1414 (August)
- Goyal, A., and I. Welch (2003). "Predicting the equity premium with dividend ratios", *Management Science* 49, 639-654.
- Hamilton J D (1989) "A New Approach to the Economic Analysis of Non-Stationary Time Series and the Business Cycle" *Econometrica* 57:357-84
- Hamilton J D (1994) *Time Series Analysis* 1994 Princeton University Press
- Kim M J C R Nelson and R Startz (1991) "Mean Reversion in Stock Prices? A Reappraisal of the Empirical Evidence" *The Review of Economic Studies*, Vol. 58, No. 3, pp. 515-528
- Lippi, M and L Reichlin (1994) "VAR analysis, nonfundamental representations, Blaschke matrices", *Journal of Econometrics*, 63 pp 307-325
- Pesaran M.H., and A. Timmerman (1995): "Predictability of Stock Returns: Robustness and Economic Significance" *Journal of Finance*, 50, 1201-1228.
- Pastor L and R F Stambaugh (2008) "Are Stocks Really Less Volatile in the Long Run?" *Working Paper*

Poterba, J.M. and L H Summers, (1988). “Mean reversion in stock prices: evidence and implications”. *Journal of Financial Economics*. 22, 27–59.

Robertson D and S Wright (2009a) “Bounds for the Predictive R-Squared” *Working Paper*

Robertson D and S Wright (2009b) “Testing for Redundant Predictor Variables” *Working Paper*

Shiller R J (2000) *Irrational Exuberance* Princeton University Press

Siegel J J (1998) *Stocks for the Long Run* McGraw-Hill

Stambaugh R F (1999) “Predictive Regressions”, *Journal of Financial Economics* 54: 375–421.

Timmermann, A, and B. Paye, 2006, “Instability of Return Prediction Models,” *Journal of Empirical Finance*, 13, 274–315.