

# To Pool or not to Pool: Revisited

M. Hashem Pesaran\*

Department of Economics, USC Dornsife INET, University of Southern California, and Trinity College, Cambridge

Qiankun Zhou

Department of Economics, Louisiana State University, Baton Rouge, LA, 70803.

October 08, 2017

## Abstract

This paper provides a new comparative analysis of pooled least squares and fixed effects estimators of the slope coefficients in the case of panel data models when the time dimension ( $T$ ) is fixed while the cross section dimension ( $N$ ) is allowed to increase without bounds. The individual effects are allowed to be correlated with the regressors, and the comparison is carried out in terms of an exponent coefficient,  $\delta$ , which measures the degree of pervasiveness of the fixed effects in the panel. The use of  $\delta$  allows us to distinguish between poolability of small  $N$  dimensional panels with large  $T$  from large  $N$  dimensional panels with small  $T$ . It is shown that the pooled estimator remains consistent so long as  $\delta < 1$ , and is asymptotically normally distributed if  $\delta < 1/2$ , for a fixed  $T$  and as  $N \rightarrow \infty$ . It is further shown that when  $\delta < 1/2$ , the pooled estimator is more efficient than the fixed effects estimator. We also propose a Hausman type diagnostic test of  $\delta < 1/2$  as a simple test of poolability, and propose a pretest estimator that could be used in practice. Monte Carlo evidence supports the main theoretical findings and gives some indications of gains to be made from pooling when  $\delta < 1/2$ .

*Keywords:* Short panel, Fixed effects estimator, Pooled estimator, Pretest estimator, Efficiency, Diagnostic test

*JEL classification:* C01, C23, C33

---

\*We would like to thank the editor, two anonymous referees, Ron Smith and Carlos Lamarche for helpful comments. Corresponding Author: Department of Economics, University of Southern California, 3620 South Vermont Avenue, Kaprielian Hall 300, Los Angeles, CA 90089, USA, pesaran@usc.edu

# 1 Introduction

This paper re-examines the issue of pooling in standard panel data models with exogenous regressors in terms of an exponent coefficient,  $0 \leq \delta \leq 1$ , which measures the degree of pervasiveness of correlated individual effects, defined by

$$\sum_{i=1}^N E |\eta_i| = O(N^\delta),$$

where  $N$  is the cross-section dimension of the panel, and  $\eta_i$  is the mean zero random part of the individual effects. The use of exponent  $\delta$  allows us to distinguish between poolability of small  $N$  dimensional panels with large  $T$  from large  $N$  dimensional panels with small  $T$ . A set of coefficients could be heterogeneous for a finite  $N$ , nevertheless can be deemed as asymptotically homogeneous if their dispersion tends to zero as  $N \rightarrow \infty$ . We use this idea to motivate conditions under which pooling is valid in large  $N$  dimensional panels, both when  $T$  is fixed and when it rises with  $N$ .

Throughout we allow for non-zero correlations between the individual effects and the regressors, and as a result the pooled estimators will be biased in the standard case where  $\delta = 1$ . We show that the choice between the pooled least squares (PLS) estimator and the fixed effects (FE) estimator depends on the value of  $\delta$ , with the PLS estimator being consistent for all values of  $\delta$  except when  $\delta = 1$ . For inference, the validity of the PLS estimator requires  $\delta < 1/2$ . Both of these conditions are significantly weaker than the homogeneity assumption made in the literature requiring that  $E |\eta_i| = 0$  for all  $i$ . For example, when  $\delta = 0$  we could have a finite number of non-zero  $E |\eta_i|$ , or more generally when  $E |\eta_i| = K\rho^i$ , for a fixed positive constant  $K$ , and  $0 < \rho < 1$ . This corresponds to the sparsity assumption often made in the context of penalized regressions. But our analysis covers non-sparse structures by allowing the number of non-zero  $E |\eta_i|$ 's to rise with  $N$  but not proportionately. The degree to which the number of units with non-zero  $E |\eta_i|$  is allowed to rise with  $N$  is governed by  $\delta$ . For example, when  $\delta = 1/2$  the number of cross-section units with non-zero random effects could rise with  $\sqrt{N}$ , with the proportion of such units in total declining to zero at the rate of  $N^{-1/2}$ .

The exponent of pervasiveness of individual effects is also closely related to the exponent of cross-sectional dependence,  $\alpha$ , recently introduced in Bailey et al. (2016) to measure the degree of cross-sectional dependence in panels. Both exponents measure the degree of pervasiveness of heterogeneity,  $\delta$  relates to the heterogeneity of the individual effects, and  $\alpha$  the heterogeneity of factor loadings in a panel data model with a factor error structure. In a broad sense,  $\delta$  can also be viewed as an exponent of cross-sectional dependence applied to the intercepts viewed as a common factor.

Our analysis complements and provides further insights on the discussion of "pool or not to pool" in the panel literature.<sup>1</sup> See for example, Baltagi et al. (2000), and Baltagi (2008). More specifically, we derive the asymptotic properties of the pooled least squares estimator when  $N$  is large and  $T$  is fixed for different values of  $\delta$ , and derive the bias of PLS when  $\delta = 1$ , and show that the pooled estimator is more efficient than the fixed effects estimator if  $\delta < 1/2$ . We also establish the asymptotic equivalence of random effects and PLS estimators when  $\delta < 1$ . In the case where  $N$  and  $T \rightarrow \infty$ , such that  $T = O(N^d)$ , for some  $d > 0$ , the condition for poolability generalizes to  $\delta < (1 - d)/2$ .

The analysis of this paper also shows the importance of knowing  $\delta$  in the choice between PLS (or RE) and FE estimators. In the case of large  $N$  and  $T$  panels estimation of  $\delta$  can be carried out using the approach of Bailey et al. (2016). But for short  $T$  panels, which is of concern in this paper, such an approach will not be applicable and other suitable techniques will be required. Accordingly, we propose a Hausman type diagnostic test of  $\delta < (1 - d)/2$  which could be used in practice as a simple test of poolability of panel data models. Finally, as an alternative strategy, we also propose a pretest estimator using a Hausman type diagnostic test and derive its asymptotic properties.

Monte Carlo simulations are conducted to compare the finite sample properties, PLS, FE and the pretest estimators. The results confirm the main theoretical findings and give some indication of the magnitudes of the gains involved from pooling when  $\delta < (1 - d)/2$ . The Monte Carlo results also place the small sample performance of the pretest estimator somewhere between those of PLS and FE estimators and is to be recommended in practice where it is not known whether  $\delta < (1 - d)/2$ .

The rest of the paper is organized as follows. Section 2 sets out the model and its assumptions. Section 3 presents the main theoretical results on the consistency and asymptotic normality of PLS and FE estimators in terms of different values of  $\delta$ . The diagnostic test of poolability is presented in Section 4. The pretest estimator is discussed in Section 5. Monte Carlo simulations are provided in Section 6, with some concluding remarks in Section 7. All mathematical derivations are provided in the Appendix.

---

<sup>1</sup>There is also a related literature that considers the problem of pooling more generally and discusses the issue of pooling in the case of panel data models with heterogenous slopes. As a recent example, see Paap, Wang and Zhang (2015) and references cited therein. In this paper we focus on the issue of pooling in the context of standard panel data models with homogeneous slopes. But our approach and generalization of the concept of cross-sectional heterogeneity can also be applied to panel data models with heterogeneous slopes.

## 2 Panel data model

Consider the standard panel data model

$$y_{it} = \alpha_i + \beta' \mathbf{x}_{it} + u_{it}, \text{ for } i = 1, 2, \dots, N; t = 1, 2, \dots, T \quad (2.1)$$

$$\alpha_i = \alpha + \eta_i \text{ for } i = 1, 2, \dots, N, \quad (2.2)$$

where  $\alpha_i$  are the individual effects,  $\mathbf{x}_{it}$  is a  $k \times 1$  vector of regressors which we decompose as

$$\mathbf{x}_{it} = \eta_i \mathbf{g}_t + \mathbf{w}_{it}, \text{ for } i = 1, 2, \dots, N; t = 1, 2, \dots, T, \quad (2.3)$$

where  $\eta_i \mathbf{g}_t$  represents the part of  $\mathbf{x}_{it}$  which is correlated with the individual effects,  $\alpha_i$ , with  $\mathbf{g}_t$  being a  $k \times 1$  vector of time effects, and  $\mathbf{w}_{it}$  is the part of  $\mathbf{x}_{it}$  which is distributed independently of the individual effects. This is a fairly general specification which allows for non-zero, possibly time-varying, correlations between  $\mathbf{x}_{it}$  and  $\alpha_i$ , and allows the regressors to have individual-specific effects and be cross-sectionally correlated. Additional individual-specific effects can be included in  $\mathbf{x}_{it}$  through  $\mathbf{w}_{it}$ . For example, using (2.3), and assuming that  $\bar{\mathbf{g}} = T^{-1} \sum_{t=1}^T \mathbf{g}_t \neq \mathbf{0}$ , then

$$\eta_i = \boldsymbol{\pi}' \bar{\mathbf{x}}_i + v_i, \quad (2.4)$$

where

$$\begin{aligned} \boldsymbol{\pi}' &= (\bar{\mathbf{g}}' \bar{\mathbf{g}})^{-1} \bar{\mathbf{g}}', \quad v_i = -(\bar{\mathbf{g}}' \bar{\mathbf{g}})^{-1} \bar{\mathbf{g}}' \bar{\mathbf{w}}_i, \\ \bar{\mathbf{x}}_i &= T^{-1} \sum_{t=1}^T \mathbf{x}_{it}, \text{ and } \bar{\mathbf{w}}_i = T^{-1} \sum_{t=1}^T \mathbf{w}_{it}, \end{aligned}$$

which is the same as Mundlak (1978) formulation of the individual effects in standard panel data models.

Throughout we assume  $T$  is fixed and carry out our analysis for  $N$  large. Except for the assumption regarding the individual effects,  $\eta_i$ , we make the following standard assumptions:

**Assumption 1:** The individual effects,  $\eta_i$  for  $i = 1, 2, \dots, N$ , are either deterministic and bounded (i.e.  $|\eta_i| < K$ ), or stochastic with second order moments,  $E(\eta_i^2) < K$ , and distributed independently of  $\mathbf{g}_t$  and  $\mathbf{w}_{jt}$  for all  $i, j$  and  $t$ ; satisfying the conditions<sup>2</sup>

$$N^{-1} \sum_{i=1}^N E|\eta_i|^s = O(N^{\delta-1}), \text{ for } s = 1 \text{ and } 2, \text{ where } 0 \leq \delta \leq 1. \quad (2.5)$$

---

<sup>2</sup> $K$  represents a generic finite positive constant.

**Remark 2.1** *The conditions of Assumption 1 are satisfied, for example, if there exists an ordering of the individual units such that for  $\delta$  in the range  $[0, 1]$*

$$\begin{aligned}\eta_i &= \varepsilon_i, \text{ for } i = 1, 2, \dots, [N^\delta], \\ &= 0, \text{ for } i = [N^\delta] + 1, [N^\delta] + 2, \dots, N\end{aligned}$$

where  $\{\varepsilon_i, i = 1, 2, \dots, N\}$  is a sequence of random variables with zero means and finite variances such that as  $M \rightarrow \infty$

$$\frac{1}{M} \sum_{i=1}^M E |\varepsilon_i|^s = O(1), \text{ for } s = 1 \text{ and } 2.$$

Then,

$$N^{-1} \sum_{i=1}^N E |\eta_i|^s = N^{\delta-1} \left( N^{-\delta} \sum_{i=1}^{[N^\delta]} |\varepsilon_i|^s \right) = O(N^{\delta-1}).$$

Note that the above result holds even if  $\varepsilon_i$ 's are cross-sectionally correlated. Furthermore, the condition that  $\eta_i = 0$ , for  $i = [N^\delta] + 1, [N^\delta] + 2, \dots, N$ , can be relaxed by requiring (See also Bailey et al. (2016)).

$$\sum_{i=[N^\delta]+1}^N E |\eta_i|^s = O(1), \text{ for } s = 1 \text{ and } 2.$$

This condition holds, for example, if  $E |\eta_i|^s = \kappa_{is} \rho_s^i$ , for  $i = [N^\delta] + 1, [N^\delta] + 2, \dots, N$ , where  $\kappa_{is}$  are finite positive constants and  $0 \leq \rho_s < 1$ .

**Remark 2.2** *Conditions (2.5) also imply*

$$N^{-1} \sum_{i=1}^N \eta_i^2 = O_p(N^{\delta-1}), \text{ and } N^{-1} \sum_{i=1}^N |\eta_i| = O_p(N^{\delta-1}).$$

These results follow by application of Markov inequality to (2.5).

**Assumption 2:** (a)  $u_{it}$  is distributed independently of  $\eta_j$  and  $\mathbf{w}_{jt'}$  for all  $i, j, t$ , and  $t'$ . (b)  $u_{it} \sim IID(0, \sigma_u^2)$  and  $E |u_{it}|^{4+\epsilon} < K < \infty$ , for some small positive  $\epsilon$ .

**Assumption 3:** The time effects,  $\mathbf{g}_t$ , are bounded such that  $\|\mathbf{g}_t \mathbf{g}_t'\| < K < \infty$ , if  $\mathbf{g}_t$  is deterministic and  $E \|\mathbf{g}_t \mathbf{g}_t'\| < K < \infty$ , if  $\mathbf{g}_t$  is stochastic.  $\|\mathbf{A}\|$  represents the Frobenius norm of  $\mathbf{A}$  defined by  $[Tr(\mathbf{A}\mathbf{A}')]^{1/2}$ .

**Assumption 4:** The variables,  $\mathbf{w}_{it}$ , are either deterministic and bounded, namely  $\|\mathbf{w}_{it}\| < K < \infty$ , or they satisfy the moment conditions  $E \|\mathbf{w}_{it} - \bar{\mathbf{w}}_i\|^2 < K < \infty$ , for all  $i$  and  $t$ , where  $\bar{\mathbf{w}}_i = T^{-1} \sum_{t=1}^T \mathbf{w}_{it}$ . Similarly,  $E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\|^2 < K < \infty$ , for all  $i$ , where  $\bar{\mathbf{w}} = N^{-1} \sum_{i=1}^N \bar{\mathbf{w}}_i$ .

**Assumption 5:** The  $k \times k$  matrices

$$\begin{aligned}\boldsymbol{\Omega}_{P,N} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})', \\ \boldsymbol{\Omega}_{FE,N} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)',\end{aligned}$$

are positive definite for all  $T$ , and as  $N \rightarrow \infty$ . The probability limits of  $\boldsymbol{\Omega}_{P,N}$  and  $\boldsymbol{\Omega}_{FE,N}$ , as  $N$  tends to infinity, will be denoted by  $\boldsymbol{\Omega}_P$  and  $\boldsymbol{\Omega}_{FE}$ , respectively.

**Remark 2.3** Note that  $E \|\mathbf{w}_{it} - \bar{\mathbf{w}}_i\| \leq \left[ E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\|^2 \right]^{1/2} < K$ , and  $E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\| \leq \left[ E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\|^2 \right]^{1/2} < K$ . Hence under Assumption 4 we also have

$$E \|(\mathbf{w}_{it} - \bar{\mathbf{w}})\| = E \|(\mathbf{w}_{it} - \bar{\mathbf{w}}_i + \bar{\mathbf{w}}_i - \bar{\mathbf{w}})\| \leq E \|(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)\| + E \|(\bar{\mathbf{w}}_i - \bar{\mathbf{w}})\| < K. \quad (2.6)$$

### 3 Pooled least squares and FE estimators

The PLS and FE estimators,  $\hat{\boldsymbol{\beta}}_P$  and  $\hat{\boldsymbol{\beta}}_{FE}$ , respectively, can be written as

$$\hat{\boldsymbol{\beta}}_P = \mathbf{Q}_{P,N}^{-1} \mathbf{q}_{P,N}, \quad (3.1)$$

and

$$\hat{\boldsymbol{\beta}}_{FE} = \mathbf{Q}_{FE,N}^{-1} \mathbf{q}_{FE,N}, \quad (3.2)$$

where

$$\mathbf{Q}_{P,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) (\mathbf{x}_{it} - \bar{\mathbf{x}})', \quad \mathbf{q}_{P,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) (y_{it} - \bar{y}), \quad (3.3)$$

$$\mathbf{Q}_{FE,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (\mathbf{x}_{it} - \bar{\mathbf{x}}_i)', \quad \mathbf{q}_{FE,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (y_{it} - \bar{y}_i), \quad (3.4)$$

and

$$\bar{\mathbf{x}} = (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T \mathbf{x}_{it}, \quad \bar{y} = (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^T y_{it}, \quad (3.5)$$

$$\bar{\mathbf{x}}_i = T^{-1} \sum_{t=1}^T \mathbf{x}_{it}, \quad \bar{y}_i = T^{-1} \sum_{t=1}^T y_{it}. \quad (3.6)$$

### 3.1 The PLS estimator

For the pooled least square estimator (PLS) defined in (3.1), we note that it can be rewritten as

$$\hat{\beta}_P = \beta + \mathbf{Q}_{P,N}^{-1} \left[ \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\eta_i + u_{it}) + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}) \right], \quad (3.7)$$

and it is shown in the appendix that under  $\delta < 1$  (for a fixed  $T$  and as  $N \rightarrow \infty$ )

$$\mathbf{Q}_{P,N} \rightarrow_p \mathbf{\Omega}_P = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E [(\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})'] > 0. \quad (3.8)$$

and

$$\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\eta_i + u_{it}) + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}) = O_p(N^{\delta-1}). \quad (3.9)$$

Substituting (3.8) and (3.9) into (3.7) now establishes the following proposition.

**Proposition 3.1** *Consider the panel data model defined by (2.1), (2.2) and (2.3). Further, suppose that Assumptions 1-5 hold. Then the PLS estimator,  $\hat{\beta}_P$ , defined by (3.1) is consistent for  $\beta$ , as long as  $\delta < 1$ .*

**Remark 3.2** *The asymptotic bias of the PLS estimator in the case of  $\delta = 1$  is given by*

$$plim_{N \rightarrow \infty} (\hat{\beta}_P) = \beta + \sigma_\eta^2 \mathbf{Q}_P^{-1} \bar{\mathbf{g}},$$

where

$$\sigma_\eta^2 = \lim_{N \rightarrow \infty} N^{-1} \sum_{i=1}^N (\eta_i - \bar{\eta})^2,$$

and

$$\mathbf{Q}_P = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E [(\mathbf{x}_{it} - \bar{\mathbf{x}}) (\mathbf{x}_{it} - \bar{\mathbf{x}})'] .$$

For a derivation see Section 26.3 in Pesaran (2015). As a corollary it also follows that Hausman's (1978) mis-specification test that compares the pooled and FE estimators will only be consistent if  $\delta = 1$ .

To derive the asymptotic distribution of  $\hat{\beta}_P$ , following the derivation in the appendix, we have

$$\sqrt{N} (\hat{\beta}_P - \beta) = \mathbf{\Omega}_P^{-1} \left[ \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) u_{it} \right] + O_p(N^{\delta-1/2}). \quad (3.10)$$

Also under Assumptions 2, 4 and 5, using standard results from panel data literature (e.g., Pesaran (2015, Chapter 26)), we have (for a fixed  $T$  and as  $N \rightarrow \infty$ )

$$\frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) u_{it} \rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \mathbf{\Omega}_P).$$

Hence, for a fixed  $T$  and as  $N \rightarrow \infty$

$$\sqrt{N} (\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}) \rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \mathbf{\Omega}_P^{-1}), \text{ if } \delta < 1/2. \quad (3.11)$$

### 3.2 The FE estimator

Consider now the FE estimator,  $\hat{\boldsymbol{\beta}}_{FE}$ , defined by (3.2). Then using (3.4) we obtain

$$\sqrt{N} (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}) = \mathbf{Q}_{FE,N}^{-1} \left[ \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (u_{it} - \bar{u}_i) \right], \quad (3.12)$$

and it is shown in the appendix that

$$\sqrt{N} (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}) = \mathbf{Q}_{FE,N}^{-1} \left[ \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (u_{it} - \bar{u}_i) \right] + O(N^{\delta-1/2}). \quad (3.13)$$

Therefore, for a fixed  $T$  and as  $N \rightarrow \infty$ , we have

$$\sqrt{N} (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}) \rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \mathbf{\Omega}_{FE}^{-1}), \text{ for } \delta < 1/2. \quad (3.14)$$

Using (3.11) and the above result now yields the following proposition:

**Proposition 3.3** *Suppose that the exponent coefficient,  $\delta$ , defined by Assumption 1, is less than 1/2, and Assumptions 1-5 hold. Then for a fixed  $T$ , and as  $N \rightarrow \infty$*

$$\sqrt{NT} (\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}) \rightarrow_d N(\mathbf{0}, \sigma_u^2 \mathbf{\Omega}_P^{-1}),$$

and

$$\sqrt{NT} (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}) \rightarrow_d N(\mathbf{0}, \sigma_u^2 \mathbf{\Omega}_{FE}^{-1}).$$

Furthermore,  $\hat{\boldsymbol{\beta}}_P$  is asymptotically more efficient than  $\hat{\boldsymbol{\beta}}_{FE}$ , as long as  $\delta < 1/2$ .

**Remark 3.4** *In the case where  $T = O(N^d)$ , for some  $d > 0$ , the condition for  $\hat{\boldsymbol{\beta}}_P$  to be asymptotically more efficient than  $\hat{\boldsymbol{\beta}}_{FE}$  is given by  $\delta < (1-d)/2$ , as  $N$  and  $T \rightarrow \infty$ . This result follows if the expressions in (3.10) and (3.13) are pre-multiplied by  $\sqrt{T}$ , and  $T$  is replaced by  $N^d$ . When  $N$  and  $T$  expand at the same rate, and  $d = 1$ , the FE estimator is always more efficient.*

For the above asymptotic distribution of PLS and FE estimators, consistent estimators of  $\mathbf{\Omega}_P$  and  $\mathbf{\Omega}_{FE}$  are given by  $\mathbf{Q}_{N,P}$  and  $\mathbf{Q}_{N,FE}$ , respectively (see (3.8)).



### 3.3 Random effects and PLS estimators

Finally, it is easily seen that random effects (RE) and the pooled least squares estimators of  $\beta$  are asymptotically equivalent. The RE estimator is given by (see, for example, Chapter 26 in Pesaran (2015)).

$$\hat{\beta}_{RE} = (\mathbf{Q}_{FE,N} + \psi \mathbf{Q}_{C,N})^{-1} (\mathbf{q}_{FE,N} + \psi \mathbf{q}_{C,N}),$$

where  $\mathbf{Q}_{FE,N}$  and  $\mathbf{q}_{FE,N}$ , are defined by (3.4),

$$\mathbf{Q}_{C,N} = N^{-1} \sum_{i=1}^N (\bar{\mathbf{x}}_i - \bar{\mathbf{x}}) (\bar{\mathbf{x}}_i - \bar{\mathbf{x}})', \quad \mathbf{q}_{C,N} = N^{-1} \sum_{i=1}^N (\bar{\mathbf{x}}_i - \bar{\mathbf{x}}) (\bar{y}_i - \bar{y}).$$

and

$$\psi = \frac{\sigma_u^2}{T\sigma_\eta^2 + \sigma_u^2}. \quad (3.15)$$

However, under (2.5),  $\sigma_\eta^2 = O(N^{\delta-1})$ , and for a fixed  $T$ , we have  $\psi = 1 + O(N^{\delta-1})$ , and using (3.3) and (3.4) we obtain<sup>3</sup>

$$\begin{aligned} \mathbf{Q}_{FE,N} + \psi \mathbf{Q}_{C,N} &= (\psi - 1) \mathbf{Q}_{C,N} + \mathbf{Q}_{P,N}, \\ \mathbf{q}_{FE,N} + \psi \mathbf{q}_{C,N} &= (\psi - 1) \mathbf{q}_{C,N} + \mathbf{q}_{P,N} \end{aligned}$$

Hence (for a fixed  $T$ )

$$\sqrt{N} \left( \hat{\beta}_{RE} - \hat{\beta}_P \right) \rightarrow_p \mathbf{0}, \text{ as } N \rightarrow \infty, \text{ if } \delta < 1,$$

which establishes the asymptotic second order equivalence of random effects and pooled least squares estimators as  $N \rightarrow \infty$ , for  $\delta < 1$  and a fixed  $T$ .

## 4 Diagnostic test of $\delta < \frac{1}{2}$

In the above analysis, we establish the asymptotic properties of PLS and FE estimators. We also compare the efficiency of PLS and FE in Proposition (3.3) and show that the PLS estimator is more efficient than the FE estimator when  $\delta < \frac{1}{2}$ . Hence, it would be desirable to use the PLS estimator for model (2.1) in terms of efficiency if  $\delta < 1/2$ . Here we propose a Hausman type diagnostic test (Hausman, (1978)) for the test of

$$H_0 : \delta = \frac{1}{2} - \epsilon, \text{ against } H_1 : \delta \geq 1/2, \quad (4.1)$$

---

<sup>3</sup>Note tha  $\mathbf{Q}_{P,N} = \mathbf{Q}_{FE,N} + \mathbf{Q}_{C,N}$ , and  $\mathbf{q}_{P,N} = \mathbf{q}_{FE,N} + \mathbf{q}_{C,N}$ .

where  $0 < \epsilon \leq \frac{1}{2}$ . Such a test will be based on the difference between the PLS and FE estimators, given by (3.1) and (3.2), respectively. We note that under  $\delta < 1$  both estimators are consistent, but under null hypothesis (4.1), (3.1) is more efficient than (3.2). Let

$$\hat{\mathbf{q}} = \hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE}, \quad (4.2)$$

then the Hausman test examines whether the PLS and FE estimators are significantly different. Then under the null hypothesis,  $H_0$ , we have

$$\text{Var}(\hat{\mathbf{q}}) = \text{Var}(\hat{\boldsymbol{\beta}}_{FE}) - \text{Var}(\hat{\boldsymbol{\beta}}_P), \quad (4.3)$$

which can be estimated as

$$\widehat{\text{Var}}(\hat{\mathbf{q}}) = \widehat{\text{Var}}(\hat{\boldsymbol{\beta}}_{FE}) - \widehat{\text{Var}}(\hat{\boldsymbol{\beta}}_P), \quad (4.4)$$

where  $\widehat{\text{Var}}(\hat{\boldsymbol{\beta}}_{FE})$  and  $\widehat{\text{Var}}(\hat{\boldsymbol{\beta}}_P)$  are the estimated covariance matrices of  $\hat{\boldsymbol{\beta}}_{FE}$  and  $\hat{\boldsymbol{\beta}}_P$  obtained under the assumption that errors,  $u_{it}$ , are serially uncorrelated and homoscedastic. Under this setting, the Hausman test statistics is given by

$$H_N = \hat{\mathbf{q}}' \left[ \widehat{\text{Var}}(\hat{\mathbf{q}}) \right]^{-1} \hat{\mathbf{q}}, \quad (4.5)$$

which is distributed as  $\chi_k^2$  for  $N$  sufficiently large, where  $k$  is the number of regressors in model (2.1).

If the Hausman test statistics (4.5) can't reject the null hypothesis  $H_0$  in (4.1), then by using the result in Proposition 3.3, it is more efficient to pool the data and use the PLS estimator.

However, it should be noted that the above test does not apply if the errors  $u_{it}$  are serially correlated or cross-sectionally heteroscedastic. In this case, we can still show that both  $\hat{\boldsymbol{\beta}}_P$  and  $\hat{\boldsymbol{\beta}}_{FE}$  are consistent, but neither is efficient. Therefore, the Hausman formula for variance of the difference doesn't apply, namely  $\text{Var}(\hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE}) \neq \text{Var}(\hat{\boldsymbol{\beta}}_{FE}) - \text{Var}(\hat{\boldsymbol{\beta}}_P)$ . But we note that

$$\hat{\mathbf{q}} = (\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}) - (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}), \quad (4.6)$$

and (using results from the previous section)

$$E(\hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE} | \mathbf{X}) = 0. \quad (4.7)$$

Also, in order to control the effects of  $\delta$  and sample sizes  $(N, T)$ , suppose now  $T = O(N^d)$ , for some  $d > 0$ . In the case where  $T$  is fixed, then  $d$  is close to zero. In view of (4.6) we have

$$\sqrt{NT} \hat{\mathbf{q}} = \sqrt{NT} (\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}) - \sqrt{NT} (\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}),$$

and using the results in (3.10) and (3.13) we obtain

$$\sqrt{NT}\hat{\mathbf{q}} = \frac{1}{\sqrt{TN}} \sum_{i=1}^N \sum_{t=1}^T \left[ \mathbf{Q}_{P,N}^{-1} (\mathbf{w}_{it} - \bar{\mathbf{w}}) - \mathbf{Q}_{FE,N}^{-1} (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) \right] u_{it} + O_p \left( N^{\delta + \frac{d}{2} - 1/2} \right), \quad (4.8)$$

where the first term will contribute to the limiting distribution, and the second will vanish if

$$\delta + \frac{d}{2} - 1/2 < 0 \quad \text{or} \quad \delta < \frac{1}{2} (1 - d). \quad (4.9)$$

This is the same as the poolability condition discussed in Remark 3.4

**Remark 4.1** *It is of interest to note that both  $\delta$  and  $d$  have significant impact on the validity of the test. From (4.8), the second term on the RHS will disappear if and only if  $\delta < \frac{1}{2} (1 - d)$ . By definition of  $d$ , we have  $d = \ln T / \ln N$ , thus we require  $\delta < \frac{1}{2} (1 - \ln T / \ln N)$ . It should be noted that  $\ln T / \ln N$  will not be a small number even if  $T$  is fixed and  $N$  is large. For instance,  $d = 0.1590$  if  $N = 1000$  and  $T = 3$ , and  $d = 0.2330$  if  $N = 1000$  and  $T = 5$ . As a result, the magnitude of  $\ln T / \ln N$  matters for the size of the Hausman type tests, a feature which is apparent from the Monte Carlo simulations reported below.*

For the implementation of the Hausman test in the general case, by direct derivations, we have

$$\text{Var} \left( \sqrt{NT}\hat{\mathbf{q}} \right) = NT \times \text{Var} \left( \hat{\beta}_P - \hat{\beta}_{FE} \right), \quad (4.10)$$

and it is shown in the appendix that

$$\text{Var} \left( \sqrt{NT}\hat{\mathbf{q}} \right) = \begin{pmatrix} \mathbf{Q}_{FE}^{-1} \mathbf{V}_{FE} \mathbf{Q}_{FE}^{-1} + \mathbf{Q}_P^{-1} \mathbf{V}_P \mathbf{Q}_P^{-1} \\ -\mathbf{Q}_{FE}^{-1} \mathbf{V}_{FEP} \mathbf{Q}_P^{-1} - \mathbf{Q}_P^{-1} \mathbf{V}_{PFE} \mathbf{Q}_{FE}^{-1} \end{pmatrix}, \quad (4.11)$$

where  $\mathbf{Q}_{FE}$  and  $\mathbf{Q}_P$  are probability limit of  $\mathbf{Q}_{FE,N}$  and  $\mathbf{Q}_{P,N}$ , respectively, and

$$\mathbf{V}_P = \lim_{N \rightarrow \infty} \mathbf{V}_{P,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it'} - \bar{\mathbf{w}})', \quad (4.12)$$

with  $\gamma_i(t, t') = E(u_{it} u_{it'} | \mathbf{X})$ ,

$$\mathbf{V}_{FE} = \lim_{N \rightarrow \infty} \mathbf{V}_{FE,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it'} - \bar{\mathbf{w}}_i)', \quad (4.13)$$

$$\mathbf{V}_{FEP} = \lim_{N \rightarrow \infty} \mathbf{V}_{FEP,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it'} - \bar{\mathbf{w}})', \quad (4.14)$$

and

$$\mathbf{V}_{PFE} = \lim_{N \rightarrow \infty} \mathbf{V}_{PFE,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it'} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)'. \quad (4.15)$$

**Remark 4.2** *It should be noted that for the general form (4.11), it can be easily verified that it reduces to the standard formula if the errors are assumed to be serially uncorrelated and homoscedastic. To see this, note that in the case of serially uncorrelated errors,  $\gamma_i(t, t') = 0$  if  $t \neq t'$ , and  $\gamma_i(t, t) = \sigma_u^2$ , we then have*

$$\begin{aligned}
\mathbf{V}_{FEP} &= \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sigma_u^2 (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}})' \\
&= \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sigma_u^2 (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) [(\mathbf{w}_{it} - \bar{\mathbf{w}}_i) + (\bar{\mathbf{w}}_i - \bar{\mathbf{w}})]' \\
&= \sigma_u^2 \mathbf{\Omega}_{FE} + \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sigma_u^2 (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\bar{\mathbf{w}}_i - \bar{\mathbf{w}})' \\
&= \sigma_u^2 \mathbf{\Omega}_{FE}.
\end{aligned}$$

Similarly,  $\mathbf{V}_{PFE} = \sigma_u^2 \mathbf{\Omega}_{FE}$ . Therefore, in this case

$$\begin{aligned}
\text{Var} \left( \hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE} | \mathbf{X} \right) &= \frac{1}{NT} \sigma_u^2 \begin{pmatrix} \mathbf{\Omega}_{FE}^{-1} \mathbf{\Omega}_{FE} \mathbf{\Omega}_{FE}^{-1} + \mathbf{\Omega}_P^{-1} \mathbf{\Omega}_P \mathbf{\Omega}_P^{-1} \\ -\mathbf{\Omega}_{FE}^{-1} \mathbf{\Omega}_{FE} \mathbf{\Omega}_P^{-1} - \mathbf{\Omega}_P^{-1} \mathbf{\Omega}_{FE} \mathbf{\Omega}_{FE}^{-1} \end{pmatrix} \\
&= \frac{1}{NT} \sigma_u^2 (\mathbf{\Omega}_{FE}^{-1} - \mathbf{\Omega}_P^{-1}) \\
&= \text{Var} \left( \hat{\boldsymbol{\beta}}_{FE} \right) - \text{Var} \left( \hat{\boldsymbol{\beta}}_P \right),
\end{aligned}$$

which accords with the Hausman's variance formula in (4.3). See also Section 26.9 in Pesaran (2015).

Given the consistent estimator of (4.11), a general Hausman test statistics of (4.1) has the form

$$\tilde{H}_N = \hat{\mathbf{q}}' \left[ \widehat{\text{Var}} \left( \hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE} \right) \right]^{-1} \hat{\mathbf{q}}, \quad (4.16)$$

which is distributed as  $\chi_k^2$  for  $N$  sufficiently large. In the general case where the errors are serially correlated and/or cross-sectionally heteroscedastic, using (4.11), under the null hypothesis,  $\text{Var} \left( \hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE} \right)$  can be consistently estimated by<sup>4</sup>

$$\widehat{\text{Var}} \left( \hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE} \right) = \frac{1}{NT} \begin{bmatrix} \mathbf{Q}_{FE,N}^{-1} \hat{\mathbf{V}}_{FE,N} \mathbf{Q}_{FE,N}^{-1} + \mathbf{Q}_{P,N}^{-1} \hat{\mathbf{V}}_{P,NT} \mathbf{Q}_{P,N}^{-1} \\ -\mathbf{Q}_{FE,N}^{-1} \hat{\mathbf{V}}_{FEP,N} \mathbf{Q}_{P,N}^{-1} - \mathbf{Q}_{P,N}^{-1} \hat{\mathbf{V}}_{PFE,N} \mathbf{Q}_{FE,N}^{-1} \end{bmatrix}, \quad (4.17)$$

where (see (3.8) and (A.10))

$$\mathbf{Q}_{FE,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (\mathbf{x}_{it} - \bar{\mathbf{x}}_i)'; \quad \mathbf{Q}_{P,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) (\mathbf{x}_{it} - \bar{\mathbf{x}})',$$

---

<sup>4</sup>See Pesaran (2015, pp 653-655).

and

$$\begin{aligned}\hat{\mathbf{V}}_{FE,N} &= \frac{1}{NT} \sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_T \hat{\mathbf{u}}_i^* \hat{\mathbf{u}}_i^{*'} \mathbf{M}_T \mathbf{X}_i, \quad \hat{\mathbf{V}}_{P,NT} = \frac{1}{NT} \sum_{i=1}^N \tilde{\mathbf{X}}'_i \hat{\mathbf{u}}_{i,OLS} \hat{\mathbf{u}}'_{i,OLS} \tilde{\mathbf{X}}_i, \\ \hat{\mathbf{V}}_{FEP,N} &= \frac{1}{NT} \sum_{i=1}^N \mathbf{X}'_i \mathbf{M}_T \hat{\mathbf{u}}_{i,OLS} \hat{\mathbf{u}}'_{i,OLS} \tilde{\mathbf{X}}_i, \quad \hat{\mathbf{V}}_{PFE,N} = \hat{\mathbf{V}}'_{FEP,N},\end{aligned}$$

where  $\mathbf{X}'_i = (\mathbf{x}_{i1}, \mathbf{x}_{i2}, \dots, \mathbf{x}_{iT})$ ,  $\hat{\mathbf{u}}_i^* = \mathbf{M}_T (\mathbf{y}_i - \mathbf{X}_i \hat{\boldsymbol{\beta}}_{FE})$ ,  $\tilde{\mathbf{X}}'_i = (\mathbf{x}_{i1} - \bar{\mathbf{x}}, \mathbf{x}_{i2} - \bar{\mathbf{x}}, \dots, \mathbf{x}_{iT} - \bar{\mathbf{x}})$ ,  $\hat{\mathbf{u}}_{i,OLS} = (\hat{u}_{i1,OLS}, \hat{u}_{i2,OLS}, \dots, \hat{u}_{iT,OLS})'$  with  $\hat{u}_{it,OLS} = y_{it} - \bar{y} - (\mathbf{x}_{it} - \bar{\mathbf{x}})' \hat{\boldsymbol{\beta}}_P$  for  $t = 1, \dots, T$ , and  $\mathbf{M}_T = \mathbf{I}_T - \boldsymbol{\tau}_T (\boldsymbol{\tau}'_T \boldsymbol{\tau}_T)^{-1} \boldsymbol{\tau}'_T$  with  $\boldsymbol{\tau}_T$  being a vector of ones of length  $T$ .

## 5 A pretest estimator

Using Hausman test statistic,  $\tilde{H}_N$ , given by (4.16), the following pretest estimator can also be considered

$$\hat{\boldsymbol{\beta}}_{Pretest} = \hat{\boldsymbol{\beta}}_P + 1 \left( \tilde{H}_N > \chi_{k,1-\tau}^2 \right) \left( \hat{\boldsymbol{\beta}}_{FE} - \hat{\boldsymbol{\beta}}_P \right), \quad (5.1)$$

where  $1(A)$  is the indicator function which takes the value of unity if  $A > 0$  and zero otherwise,  $\tau$  is the nominal level of the chi-square distribution with  $k$  degrees of freedom ( $\chi_k^2$ ). Following the line of proof in Guggenberger (2010), Kabaila et al. (2015) and Wang et al. (2016), the asymptotic distribution of  $\hat{\boldsymbol{\beta}}_{Pretest}$  is established in the following proposition, and the proof is provided in the Appendix.

**Proposition 5.1** *Suppose that the exponent coefficient,  $\delta$ , defined by Assumption 1, is less than  $1/2$ , and Assumptions 1-5 hold. Then for a fixed  $T$ , and as  $N \rightarrow \infty$*

$$(i) \sqrt{N} \begin{pmatrix} \hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta} \\ \hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta} \end{pmatrix} \rightarrow_d \boldsymbol{\psi} = \begin{pmatrix} \boldsymbol{\psi}_1 \\ \boldsymbol{\psi}_2 \end{pmatrix} \sim N(0, \mathbf{V}), \text{ with } \mathbf{V} = \begin{pmatrix} \mathbf{V}_P & \mathbf{V}_P \\ \mathbf{V}_P & \mathbf{V}_{FE} \end{pmatrix},$$

where  $\mathbf{V}_P = \sigma_u^2 T^{-1} \boldsymbol{\Omega}_P^{-1}$ , and  $\mathbf{V}_{FE} = \sigma_u^2 T^{-1} \boldsymbol{\Omega}_{FE}^{-1}$ ;

$$(ii) \tilde{H}_N \rightarrow_d \varphi = \boldsymbol{\psi}'_3 \boldsymbol{\psi}_3, \text{ where } \boldsymbol{\psi}_3 = \boldsymbol{\Omega} (\boldsymbol{\psi}_2 - \boldsymbol{\psi}_1) \sim N(\mathbf{0}, \mathbf{I}_k), \text{ with } \boldsymbol{\Omega} = (\mathbf{V}_{FE} - \mathbf{V}_P)^{-1/2};$$

$$(iii) \sqrt{N} \left( \hat{\boldsymbol{\beta}}_{Pretest} - \boldsymbol{\beta} \right) \rightarrow_d \boldsymbol{\psi}_1 + \boldsymbol{\Omega}^{-1} \boldsymbol{\psi}_3 1 \left( \varphi > \chi_{k,1-\tau}^2 \right).$$

**Remark 5.2** *The above results can be easily extend to the case where  $T = O(N^d)$ , with  $d > 0$ , i.e.,  $T \rightarrow \infty$  as  $N \rightarrow \infty$ . For this scenario, we note that as  $(N, T) \rightarrow \infty$ , we have*

$$\sqrt{NT} \left( \hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta} \right) \rightarrow_d N(\mathbf{0}, \sigma_u^2 \boldsymbol{\Omega}_P^{-1}),$$

and

$$\sqrt{NT} \left( \hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta} \right) \rightarrow_d N(\mathbf{0}, \sigma_u^2 \boldsymbol{\Omega}_{FE}^{-1}),$$

Furthermore,  $\hat{\beta}_P$  is asymptotically more efficient than  $\hat{\beta}_{FE}$ , as long as  $\delta < 1/2$ . For the hypothesis testing problem (4.1), the test statistic is given by (4.5) and the pretest estimator is defined by (5.1). The limiting distribution of (5.1) is given by

$$(i) \sqrt{NT} \begin{pmatrix} \hat{\beta}_P - \beta \\ \hat{\beta}_{FE} - \beta \end{pmatrix} \rightarrow_d \psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \sim N(\mathbf{0}, \mathbf{V}) \text{ with } \mathbf{V} = \begin{pmatrix} \mathbf{V}_P & \mathbf{V}_P \\ \mathbf{V}_P & \mathbf{V}_{FE} \end{pmatrix},$$

where  $\mathbf{V}_P = \sigma_u^2 \mathbf{\Omega}_P^{-1}$ , and  $\mathbf{V}_{FE} = \sigma_u^2 \mathbf{\Omega}_{FE}^{-1}$ ;

$$(ii) \tilde{H}_N \rightarrow_d \varphi = \psi_3' \psi_3, \text{ where } \psi_3 = \mathbf{\Omega}(\psi_2 - \psi_1) \sim N(\mathbf{0}, \mathbf{I}_k), \text{ with } \mathbf{\Omega} = (\mathbf{V}_{FE} - \mathbf{V}_P)^{-1/2};$$

$$(iii) \sqrt{NT} (\hat{\beta}_{Pretest} - \beta) \rightarrow_d \psi_1 + \mathbf{\Omega}^{-1} \psi_3 1(\varphi > \chi_{k,1-\tau}^2);$$

## 6 Monte Carlo simulations

To compare the performance of the FE and pooled least square estimators when  $T$  is fixed as well as  $\sum_{i=1}^N |\eta_i| = O(N^\delta)$ , we conduct several Monte Carlo simulations. The data generating process (DGP) is given by

$$y_{it} = 1 + \eta_i + x_{1,it}\beta_1 + x_{2,it}\beta_2 + u_{it}, \quad i = 1, 2, \dots, N; t = 1, 2, \dots, T, \quad (6.1)$$

with  $\beta_1 = 1$  and  $\beta_2 = 2$ ,  $N = 100, 500, 1000, 2000$  and  $T = 5$ .<sup>5</sup> We assume  $u_{it} \sim IIDN(0, \sigma_i^2)$ , with  $\sigma_i^2 \sim IID\chi^2(2)$ ,  $\eta_i \sim IIDN(0, 2)$  for  $i = 1, 2, \dots, [N^\delta]$  and  $\eta_i = 0$ , for  $i = [N^\delta] + 1, [N^\delta] + 2, \dots, N$ . We let  $\delta$  to take the following values 1, 0.75, 0.5, 0.48, 0.46, 0.44, 0.42, 0.4, 0.25 and 0. The elements of  $\mathbf{x}_{it} = (x_{1,it}, x_{2,it})'$ , are generated as

$$x_{j,it} = 1 + \alpha_{j,i} + g_{j,t}\eta_i + w_{j,it}, \text{ for } j = 1, 2,$$

with  $\alpha_{j,i} \sim IIDN(0, 1)$ ,  $g_{j,t} \sim IIDU[0.1, 0.9]$ , and  $w_{j,it}$  generated by

$$w_{j,it} = \rho_{j,i} w_{j,it-1} + \varepsilon_{j,it}, \text{ for } j = 1, 2,$$

where  $w_{j,i0} = 0$ ,  $\rho_{j,i} \sim IIDU[0.05, 0.95]$ ,  $\varepsilon_{j,i0} = 0$ , and  $\varepsilon_{j,it} \sim IIDN(0, \sigma_{j,\varepsilon i}^2)$  with  $\sigma_{j,\varepsilon i}^2 \sim IID\chi^2(2)$  for  $j = 1, 2$ . For the DGP described above, the first 50 observations are discarded, and the number of replications is set to 1000.

We compute the PLS, FE and Pretest estimates and the associated bias, absolute bias and RMSE. These estimation results are summarized in Table 1-4. In line with our theoretical results, we note that the RMSE of the PLS estimator is much smaller than those of the FE estimator for values of  $\delta < 1/2$ . However, the PLS estimator starts to show significant bias as

<sup>5</sup>To save space, here we only provide the simulation results of  $T = 5$ . The simulation results of  $T = 3, 10$  are provided in the appendix.

$\delta$  is allowed to increase beyond the  $1/2$  threshold, and the RMSE of the PLS estimator is much larger than the FE estimator.

As far as the Pretest estimator is concerned, again as to be expected, we observe from the simulation results that it performs better than the FE estimator when  $\delta < 1/2$ , but is less efficient than the PLS estimator. However, when  $\delta > 1/2$ , the pretest estimator is more efficient than the PLS estimator, but not as efficient as the FE estimator.

The results of the poolability test using the Hausman type statistic are summarized in Table 5. We can observe that the empirical size is very close to the 5% nominal value when  $\delta < 0.25$ , which makes sense since for all our combinations of  $(N, T)$ , the minimum  $d$  is 0.1445 and the maximum  $d$  is 0.5. The proposed poolability test has good power for values of  $\delta > (1 - d)/2$ , as predicted by the theory.

## 7 Conclusion

This paper introduces a new approach to the analysis of the relative efficiency of fixed effects and pooled least square estimators for standard panel data models. We show that the potential benefit from pooling is directly related to the degree with which the heterogeneity of individual effects is pervasive across the individual units in the panel. We characterize this feature by an exponent,  $\delta$ , and show that pooled least square estimator is consistent for values of  $\delta < 1$ . Our specification allows for non-zero correlations between the individual effects and the regressors which renders the pooled least squares and random effects inconsistent if  $\delta = 1$ . We also derive the asymptotic distributions of the pooled least squares, FE and RE estimators for different values of  $\delta$  and establish the relative efficiency of the pooled least squares estimator over the FE estimator when  $\delta < (1 - d)/2$ , where  $d$  is given by  $\ln T / \ln N$ . We also propose a Hausman type diagnostic test of poolability, and an associated pretest estimator. The theoretical results are supported by small sample evidence from Monte Carlo experiments.

## References

- [1] Bailey, N., G. Kapetanios, and M.H. Pesaran, 2016, Exponent of cross-sectional dependence: estimation and inference, *Journal of Applied Econometrics* 31, 929-960.
- [2] Baltagi, B., G. Bresson and A. Pirotte, 2008, To Pool or Not to Pool?, Chapter 16 in László Mátyás, Patrick Sevestre (eds.), *The Econometrics of Panel Data*, 517-546, Springer Berlin Heidelberg;

- [3] Baltagi, B., J. Griffin, and W. Xiong, 2000, To pool or not to pool: homogeneous versus heterogeneous estimations applied to cigarette demand, *The Review of Economics and Statistics* 82, 117-126.
- [4] Guggenberger, P., 2010, The impact of a Hausman pretest on the size of a hypothesis test: The panel data case, *Journal of Econometrics* 156, 337-343.
- [5] Hausman, J. A. 1978. Specification tests in econometrics. *Econometrica* 46, 1251–1272.
- [6] Kabaila, P., R. Mainzer, and D. Farchione, 2015, The impact of a Hausman pretest, applied to panel data, on the coverage probability of confidence intervals, *Economics Letters* 131, 12-15,
- [7] Mundlak, Y. 1978. On the pooling of time series and cross section data. *Econometrica* 46, 69–85.
- [8] Paap, R, W. Wang, and X. Zhang 2015. To pool or not to pool: What is a good strategy? working paper.
- [9] Pesaran, M. H., 2015, *Time Series and Panel Data Econometrics*, Oxford University Press, Oxford.
- [10] Wang, Y., Y. Zhang, and Q. Zhou, 2016, A Stein-like estimator for linear panel data models, *Economics Letters* 141, 156-161.



Table 1: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 5$  and  $0 \leq \delta \leq 0.44$

$N$		$\delta$																			
		0				0.25				0.4				0.42				0.44			
		FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest		
100	Estim	0.9984	1.0012	1.0019	0.9979	1.0207	1.0169	0.9990	1.0399	1.0287	1.0002	1.0480	1.0318	0.9974	1.0463	1.0280	0.9974	1.0463	1.0280		
	Bias	-0.0016	0.0012	0.0019	-0.0021	0.0207	0.0169	-0.0010	0.0399	0.0287	0.0002	0.0480	0.0318	-0.0026	0.0463	0.0280	-0.0026	0.0463	0.0280		
	aBias	0.0483	0.0259	0.0289	0.0483	0.0281	0.0322	0.0496	0.0417	0.0463	0.0464	0.0490	0.0513	0.0457	0.0474	0.0504	0.0457	0.0474	0.0504		
	rmse	0.0606	0.0324	0.0383	0.0607	0.0348	0.0425	0.0617	0.0485	0.0557	0.0583	0.0562	0.0596	0.0575	0.0538	0.0590	0.0575	0.0538	0.0590		
500	Estim	1.0006	1.0001	0.9998	0.9982	1.0076	1.0059	1.0004	1.0193	1.0146	0.9998	1.0228	1.0148	0.9999	1.0194	1.0133	0.9999	1.0194	1.0133		
	Bias	0.0006	0.0001	-0.0002	-0.0018	0.0076	0.0059	0.0004	0.0193	0.0146	-0.0002	0.0228	0.0148	-0.0001	0.0194	0.0133	-0.0001	0.0194	0.0133		
	aBias	0.0219	0.0100	0.0121	0.0201	0.0114	0.0131	0.0206	0.0199	0.0212	0.0208	0.0230	0.0246	0.0210	0.0198	0.0216	0.0210	0.0198	0.0216		
	rmse	0.0276	0.0121	0.0174	0.0253	0.0142	0.0177	0.0258	0.0230	0.0250	0.0260	0.0256	0.0279	0.0265	0.0228	0.0228	0.0265	0.0228	0.0228		
1000	Estim	1.0002	1.0001	1.0000	0.9992	1.0057	1.0045	0.9996	1.0117	1.0088	0.9993	1.0146	1.0105	0.9988	1.0163	1.0098	0.9988	1.0163	1.0098		
	Bias	0.0002	0.0001	0.0000	-0.0008	0.0057	0.0045	-0.0004	0.0117	0.0088	-0.0007	0.0146	0.0105	-0.0012	0.0163	0.0098	-0.0012	0.0163	0.0098		
	aBias	0.0159	0.0070	0.0082	0.0152	0.0084	0.0095	0.0146	0.0125	0.0137	0.0149	0.0149	0.0161	0.0153	0.0165	0.0175	0.0153	0.0165	0.0175		
	rmse	0.0197	0.0087	0.0116	0.0188	0.0095	0.0126	0.0185	0.0147	0.0168	0.0188	0.0170	0.0189	0.0192	0.0184	0.0202	0.0192	0.0184	0.0202		
2000	Estim	0.9992	1.0001	0.9998	0.9995	1.0039	1.0030	0.9998	1.0063	1.0046	0.9999	1.0078	1.0057	0.9991	1.0109	1.0060	0.9991	1.0109	1.0060		
	Bias	-0.0008	0.0001	-0.0002	-0.0005	0.0039	0.0030	-0.0002	0.0063	0.0046	-0.0001	0.0078	0.0057	-0.0009	0.0109	0.0060	-0.0009	0.0109	0.0060		
	aBias	0.0108	0.0050	0.0061	0.0108	0.0060	0.0069	0.0106	0.0073	0.0084	0.0105	0.0083	0.0092	0.0107	0.0111	0.0121	0.0107	0.0111	0.0121		
	rmse	0.0136	0.0063	0.0087	0.0135	0.0073	0.0094	0.0134	0.0088	0.0108	0.0133	0.0092	0.0114	0.0135	0.0125	0.0141	0.0135	0.0125	0.0141		

Notes: 1. "Bias", "aBias" and "rmse" refer to bias, absolute bias and root mean square error of estimating  $\beta_1$ , respectively.

2. "FE", "PLS" and "Pretest" refer to fixed effects, pooled LS and pretest estimators defined by (3.2), (3.1) and (5.1), respectively.

Table 2: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 5$  and  $0.46 \leq \delta \leq 1$

$N$	$\delta$																			
	0.46				0.48				0.5				0.75				1			
	FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest	
100	Estim	0.9974	1.0463	1.0280	0.9983	1.0523	1.0329		1.0006	1.0607	1.0348		1.0005	1.1000	1.0075		1.0015	1.1057	1.0015	
	Bias	-0.0026	0.0463	0.0280	-0.0017	0.0523	0.0329		0.0006	0.0607	0.0348		0.0005	0.1000	0.0075		0.0015	0.1057	0.0015	
	aBias	0.0457	0.0474	0.0.504	0.0475	0.0523	0.0528		0.0491	0.0607	0.0627		0.0425	0.1000	0.0488		0.0387	0.1057	0.0387	
	rmse	0.0575	0.0538	0.0590	0.0607	0.0591	0.0641		0.0620	0.0667	0.0709		0.0537	0.1034	0.0612		0.0488	0.1087	0.0488	
500	Estim	0.9985	1.0188	1.0121	1.0001	1.0227	1.0150		0.9998	1.0295	1.0162		0.9984	1.0736	0.9985		1.0000	1.1121	1.0000	
	Bias	-0.0015	0.0188	0.0121	0.0001	0.0227	0.0150		-0.0002	0.0295	0.0162		-0.0016	0.0736	-0.0015		0.0000	0.1121	0.0000	
	aBias	0.0210	0.0193	0.0215	0.0209	0.0230	0.0243		0.0200	0.0296	0.0281		0.0197	0.0736	0.0197		0.0167	0.1121	0.0167	
	rmse	0.0263	0.0223	0.0257	0.0264	0.0243	0.0259		0.0248	0.0320	0.0311		0.0248	0.0744	0.0249		0.0207	0.1126	0.0207	
1000	Estim	0.9991	1.0161	1.0095	0.9992	1.0172	1.0096		0.9992	1.0208	1.0109		1.0000	1.0645	1.0000		1.0002	1.1132	1.0002	
	Bias	-0.0009	0.0161	0.0095	-0.0008	0.0172	0.0096		-0.0008	0.0208	0.0109		0.0000	0.0645	0.0000		0.0002	0.1132	0.0002	
	aBias	0.0145	0.0163	0.0169	0.0145	0.0174	0.0179		0.0155	0.0209	0.0210		0.0134	0.0645	0.0134		0.0124	0.1132	0.0124	
	rmse	0.0179	0.0185	0.0196	0.0183	0.0194	0.0205		0.0192	0.0226	0.0232		0.0166	0.0650	0.0166		0.0155	0.1134	0.0155	
2000	Estim	0.9999	1.0113	1.0072	0.9989	1.0132	1.0068		0.9990	1.0134	1.0071		0.9999	1.0546	0.9999		1.0001	1.1132	1.0001	
	Bias	-0.0001	0.0113	0.0072	-0.0011	0.0132	0.0068		-0.0010	0.0134	0.0071		-0.0001	0.0546	-0.0001		0.0001	0.1132	0.0001	
	aBias	0.0107	0.0114	0.0120	0.0105	0.0133	0.0137		0.0100	0.0134	0.0135		0.0095	0.0546	0.0095		0.0084	0.1132	0.0084	
	rmse	0.0133	0.0129	0.0139	0.0133	0.0145	0.0154		0.0127	0.0147	0.0151		0.0122	0.0549	0.0122		0.0105	0.1133	0.0105	

See notes of Table 1.

Table 3: Simulation results of for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 5$  and  $0 \leq \delta \leq 0.44$

N		$\delta$																				
		0				0.25				0.4				0.42				0.44				
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		
100	Estim	2.0007	2.0000	1.9994	2.0013	2.0142	2.0117	1.9999	2.0464	2.0318	2.0012	2.0448	2.0296	2.0011	2.0525	2.0333	2.0011	2.0525	2.0333	2.0011	2.0525	2.0333
	Bias	0.0007	0.0000	-0.0006	0.0013	0.0142	0.0117	-0.0001	0.0464	0.0318	0.0012	0.0448	0.0296	0.0011	0.0525	0.0333	0.0011	0.0525	0.0333	0.0011	0.0525	0.0333
	aBias	0.0480	0.0225	0.0274	0.0465	0.0242	0.0287	0.0453	0.0469	0.0506	0.0458	0.0454	0.0495	0.0444	0.0527	0.0533	0.0444	0.0527	0.0533	0.0444	0.0527	0.0533
	rmse	0.0616	0.0280	0.0390	0.0582	0.0303	0.0396	0.0567	0.0521	0.0580	0.0577	0.0510	0.0572	0.0555	0.0575	0.0600	0.0555	0.0575	0.0600	0.0555	0.0575	0.0600
500	Estim	1.9984	2.0001	1.9997	1.9986	2.0123	2.0096	1.9993	2.0175	2.0123	1.9987	2.0203	2.0133	1.9997	2.0189	2.0129	1.9997	2.0189	2.0129	1.9997	2.0189	2.0129
	Bias	-0.0016	0.0001	-0.0003	-0.0014	0.0123	0.0096	-0.0007	0.0175	0.0123	-0.0013	0.0203	0.0133	-0.0003	0.0189	0.0129	-0.0003	0.0189	0.0129	-0.0003	0.0189	0.0129
	aBias	0.0226	0.0100	0.0124	0.0219	0.0145	0.0163	0.0222	0.0184	0.0211	0.0211	0.0208	0.0228	0.0218	0.0195	0.0222	0.0218	0.0195	0.0222	0.0218	0.0195	0.0222
	rmse	0.0281	0.0128	0.0179	0.0275	0.0178	0.0210	0.0281	0.0215	0.0257	0.0266	0.0233	0.0233	0.0272	0.0222	0.0266	0.0272	0.0222	0.0266	0.0272	0.0222	0.0266
1000	Estim	1.9989	1.9995	1.9995	1.9995	2.0056	2.0042	2.0004	2.0111	2.0085	1.9999	2.0112	2.0078	2.0001	2.0164	2.0107	2.0001	2.0164	2.0107	2.0001	2.0164	2.0107
	Bias	-0.0011	-0.0005	-0.0005	-0.0005	0.0056	0.0042	0.0004	0.0111	0.0085	-0.0001	0.0112	0.0078	0.0001	0.0164	0.0107	0.0001	0.0164	0.0107	0.0001	0.0164	0.0107
	aBias	0.0158	0.0073	0.0085	0.0155	0.0084	0.0094	0.0155	0.0119	0.0132	0.0163	0.0121	0.0136	0.0161	0.0165	0.0177	0.0161	0.0165	0.0177	0.0161	0.0165	0.0177
	rmse	0.0199	0.0085	0.0120	0.0194	0.0106	0.0125	0.0194	0.0141	0.0162	0.0202	0.0143	0.0167	0.0199	0.0185	0.0207	0.0199	0.0185	0.0207	0.0199	0.0185	0.0207
2000	Estim	1.9998	1.9998	2.0000	1.9995	2.0037	2.0027	1.9998	2.0077	2.0059	1.9996	2.0079	2.0059	1.9996	2.0115	2.0066	1.9996	2.0115	2.0066	1.9996	2.0115	2.0066
	Bias	-0.0002	-0.0002	0.0000	-0.0005	0.0037	0.0027	-0.0002	0.0077	0.0059	-0.0004	0.0079	0.0059	-0.0004	0.0115	0.0066	-0.0004	0.0115	0.0066	-0.0004	0.0115	0.0066
	aBias	0.0110	0.0050	0.0058	0.0103	0.0057	0.0068	0.0112	0.0084	0.0095	0.0101	0.0085	0.0094	0.0106	0.0116	0.0122	0.0106	0.0116	0.0122	0.0106	0.0116	0.0122
	rmse	0.0136	0.0063	0.0082	0.0132	0.0068	0.0093	0.0140	0.0100	0.0118	0.0127	0.0100	0.0115	0.0134	0.0130	0.0145	0.0134	0.0130	0.0145	0.0134	0.0130	0.0145

See notes of Table 1.

Table 4: Simulation results for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 5$  and  $0.46 \leq \delta \leq 1$

N		$\delta$																					
		0.46				0.48				0.5				0.75				1					
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest			
100	Estim	2.0011	2.0525	2.0339	2.0010	2.0523	2.0336		2.0003	2.0514	2.0293		2.0016	2.0765	2.0070		1.9992	2.1244	1.9992		1.9992	2.1244	1.9992
	Bias	0.0011	0.0525	0.0339	0.0010	0.0523	0.0336		0.0003	0.0514	0.0293		0.0016	0.0765	0.0070		-0.0008	0.1244	-0.0008		-0.0008	0.1244	-0.0008
	aBias	0.0444	0.0533	0.0533	0.0462	0.0582	0.0552		0.0440	0.0519	0.0531		0.0407	0.0765	0.0454		0.0385	0.1244	0.0385		0.0385	0.1244	0.0385
	rmse	0.0555	0.0601	0.0601	0.0582	0.0583	0.0622		0.0547	0.0578	0.0609		0.0506	0.0804	0.0555		0.0476	0.1271	0.0476		0.0476	0.1271	0.0476
500	Estim	1.9996	2.0187	2.0123	2.0003	2.0264	2.0170		1.9986	2.0271	2.0140		1.9996	2.0716	1.9996		2.0006	2.1110	2.0006		2.0006	2.1110	2.0006
	Bias	-0.0004	0.0187	0.0123	0.0003	0.0264	0.0170		-0.0014	0.0271	0.0140		-0.0004	0.0716	-0.0004		0.0006	0.1110	0.0006		0.0006	0.1110	0.0006
	aBias	0.0229	0.0194	0.0221	0.0209	0.0264	0.0273		0.0203	0.0273	0.0270		0.0204	0.0716	0.0204		0.0166	0.1110	0.0166		0.0166	0.1110	0.0166
	rmse	0.0284	0.0223	0.0262	0.0263	0.0288	0.0288		0.0254	0.0298	0.0304		0.0252	0.0726	0.0253		0.0206	0.1116	0.0206		0.0206	0.1116	0.0206
1000	Estim	1.9995	2.0198	2.0117	1.9995	2.0202	2.0107		2.0002	2.0200	2.0115		2.0005	2.0618	2.0005		1.9999	2.1094	1.9999		1.9999	2.1094	1.9999
	Bias	-0.0005	0.0198	0.0117	-0.0005	0.0202	0.0107		0.0002	0.0200	0.0115		0.0005	0.0618	0.0005		-0.0001	0.1094	-0.0001		-0.0001	0.1094	-0.0001
	aBias	0.0157	0.0199	0.0202	0.0155	0.0203	0.0204		0.0154	0.0201	0.0200		0.0138	0.0618	0.0138		0.0120	0.1094	0.0120		0.0120	0.1094	0.0120
	rmse	0.0196	0.0216	0.0216	0.0194	0.0220	0.0228		0.0194	0.0221	0.0226		0.0174	0.0624	0.0174		0.0153	0.1097	0.0153		0.0153	0.1097	0.0153
2000	Estim	2.0007	2.0139	2.0088	2.0001	2.0135	2.0072		1.9993	2.0133	2.0070		2.0004	2.0558	2.0004		1.9996	2.1102	1.9996		1.9996	2.1102	1.9996
	Bias	0.0007	0.0139	0.0088	0.0001	0.0135	0.0072		-0.0007	0.0133	0.0070		0.0004	0.0558	0.0004		-0.0004	0.1102	-0.0004		-0.0004	0.1102	-0.0004
	aBias	0.0105	0.0139	0.0140	0.0108	0.0136	0.0136		0.0110	0.0138	0.0140		0.0097	0.0558	0.0097		0.0085	0.1102	0.0085		0.0085	0.1102	0.0085
	rmse	0.0133	0.0152	0.0156	0.0135	0.0149	0.0153		0.0138	0.0148	0.0158		0.0122	0.0561	0.0122		0.0106	0.1104	0.0106		0.0106	0.1104	0.0106

See notes of Table 1.

Table 5: Empirical rejection frequencies for the Hausman type test at 5% significance level

$N$	$T$	$\delta$									
		0	0.25	0.4	0.42	0.44	0.46	0.48	0.5	0.75	1
100	3	6.1%	7.4%	7.6%	6.5%	8.5%	8.5%	7.2%	8.7%	22.5%	90.6%
	5	7.6%	5.7%	8.2%	6.8%	8.1%	8.1%	8.3%	10.2%	56.7%	100%
	10	5.2%	6.7%	14.2%	9%	12.2%	12.2%	15.6%	20.7%	98.5%	100%
500	3	5.6%	5.2%	7.5%	6.2%	8.7%	7.1%	6.8%	8.5%	59.8%	100%
	5	5.9%	7%	7.4%	11.7%	12.3%	14%	18%	13.2%	99.2%	100%
	10	4.6%	5.6%	11.3%	24.9%	29.4%	43.2%	50.4%	36.9%	100%	100%
1000	3	7%	6%	6%	6.4%	6.6%	6.2%	6.8%	8.1%	71.3%	100%
	5	7.1%	5.9%	8.1%	9.7%	11.8%	12.9%	15%	14.5%	100%	100%
	10	5.2%	6.2%	21.9%	31.5%	32.6%	49.3%	58.2%	53.9%	100%	100%
2000	3	4.2%	5.4%	5.9%	5.9%	5.6%	7.3%	7.6%	9.2%	88.7%	100%
	5	4.5%	6%	6.9%	6.2%	8.5%	17.2%	15.3%	18.4%	100%	100%
	10	5.2%	4.7%	20.2%	35%	36.8%	61.5%	62.4%	71.7%	100%	100%

Note: The 5% significance level for  $\chi^2(2)$  is 5.991.

## Appendix: Mathematical derivations and additional simulation results

This appendix contains the mathematical derivations of the main theoretical results in the main paper and some additional simulation results.

### A.1 Proof of Proposition 3.1

To derive the asymptotic properties of PLS, using (2.3), we first note that

$$\mathbf{x}_{it} - \bar{\mathbf{x}} = (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) + (\mathbf{w}_{it} - \bar{\mathbf{w}}), \quad (\text{A.1})$$

and

$$y_{it} - \bar{y} = \eta_i - \bar{\eta} + \beta' (\mathbf{x}_{it} - \bar{\mathbf{x}}) + (u_{it} - \bar{u}), \quad (\text{A.2})$$

where

$$\bar{\mathbf{g}} = T^{-1} \sum_{t=1}^T \mathbf{g}_t, \quad \bar{\eta} = N^{-1} \sum_{i=1}^N \eta_i, \quad \text{and} \quad \bar{u} = N^{-1} \sum_{i=1}^N u_i.$$

Then for PLS estimator (3.1), using (A.1) in (3.3) we have

$$\begin{aligned} \mathbf{Q}_{P,N} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t + \mathbf{w}_{it} - \bar{\eta} \bar{\mathbf{g}} - \bar{\mathbf{w}}) (\eta_i \mathbf{g}_t + \mathbf{w}_{it} - \bar{\eta} \bar{\mathbf{g}} - \bar{\mathbf{w}})' \\ &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})' + \left( N^{-1} \sum_{i=1}^N \eta_i^2 \right) \left( T^{-1} \sum_{t=1}^T \mathbf{g}_t \mathbf{g}_t' \right) - \bar{\eta}^2 (\bar{\mathbf{g}} \bar{\mathbf{g}}') \\ &\quad + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i [(\mathbf{w}_{it} - \bar{\mathbf{w}}) \mathbf{g}_t' + \mathbf{g}_t (\mathbf{w}_{it} - \bar{\mathbf{w}})']. \end{aligned} \quad (\text{A.3})$$

Similarly, using (A.2) in (3.3) we have

$$\begin{aligned} \mathbf{q}_{P,N} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) (y_{it} - \bar{y}) \\ &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) [\eta_i - \bar{\eta} + \beta' (\mathbf{x}_{it} - \bar{\mathbf{x}}) + u_{it} - \bar{u}] \\ &= \mathbf{Q}_{P,N} \beta + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}) (\eta_i - \bar{\eta} + u_{it} - \bar{u}), \end{aligned}$$

which upon using (A.1) can be written as

$$\mathbf{q}_{P,N} = \mathbf{Q}_{P,N} \beta + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\eta_i + u_{it}) + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}), \quad (\text{A.4})$$

which in turn yields

$$\hat{\beta}_P = \beta + \mathbf{Q}_{P,N}^{-1} \left[ \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\eta_i + u_{it}) + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}) \right].$$

Furthermore,

$$\begin{aligned}
\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}) &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i^2 \mathbf{g}_t - \bar{\eta} \eta_i \bar{\mathbf{g}}) + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) u_{it} \\
&= \frac{1}{N} \sum_{i=1}^N (\eta_i^2 - \bar{\eta} \eta_i) \bar{\mathbf{g}} + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) u_{it} \\
&= \left[ N^{-1} \sum_{i=1}^N (\eta_i - \bar{\eta})^2 \right] \bar{\mathbf{g}} + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i \mathbf{g}_t u_{it} - \bar{\eta} \bar{\mathbf{g}} \bar{u}. \tag{A.5}
\end{aligned}$$

But under Assumption 1 we have

$$N^{-1} \sum_{i=1}^N E |\eta_i|^2 = O(N^{\delta-1}), \text{ and } E |\bar{\eta}| \leq N^{-1} \sum_{i=1}^N |\eta_i| = O(N^{\delta-1}), \tag{A.6}$$

and since  $\eta_i$  is distributed independently of  $\bar{\mathbf{g}}$  and  $u_{it}$ , then

$$\begin{aligned}
E \left| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\eta_i \mathbf{g}_t - \bar{\eta} \bar{\mathbf{g}}) (\eta_i + u_{it}) \right| &\leq \left[ N^{-1} \sum_{i=1}^N E (\eta_i - \bar{\eta})^2 \right] E (\|\bar{\mathbf{g}}\|) \\
&\quad + \sup_{i,t} E |u_{it}| \sup_t E (\|\mathbf{g}_t\|) \left( N^{-1} \sum_{i=1}^N E |\eta_i| \right) + E |\bar{\eta}| E |\bar{u}| E (\|\bar{\mathbf{g}}\|) \\
&= O(N^{\delta-1}).
\end{aligned}$$

Similarly

$$\begin{aligned}
\left\| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i [(\mathbf{w}_{it} - \bar{\mathbf{w}}) \mathbf{g}_t' + \mathbf{g}_t (\mathbf{w}_{it} - \bar{\mathbf{w}})'] \right\| &\leq \frac{2}{NT} \sum_{i=1}^N \sum_{t=1}^T |\eta_i| \|(\mathbf{w}_{it} - \bar{\mathbf{w}})\| \|\mathbf{g}_t\| \\
&= \frac{2}{N} \sum_{i=1}^N |\eta_i| \left[ T^{-1} \sum_{t=1}^T \|(\mathbf{w}_{it} - \bar{\mathbf{w}})\| \|\mathbf{g}_t\| \right],
\end{aligned}$$

and since under Assumption 1,  $\eta_i$  is distributed independently of  $\mathbf{g}_t$  and  $\mathbf{w}_{it}$ , we have

$$E \left\| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i [(\mathbf{w}_{it} - \bar{\mathbf{w}}) \mathbf{g}_t' + \mathbf{g}_t (\mathbf{w}_{it} - \bar{\mathbf{w}})'] \right\| \leq \frac{2}{N} \sum_{i=1}^N E |\eta_i| \left[ \left\{ T^{-1} \sum_{t=1}^T E [\|(\mathbf{w}_{it} - \bar{\mathbf{w}})\| \|\mathbf{g}_t\|] \right\} \right].$$

However, by Cauchy–Schwarz inequality and under Assumptions 3 and 4

$$E [\|(\mathbf{w}_{it} - \bar{\mathbf{w}})\| \|\mathbf{g}_t\|] \leq \left[ E \|(\mathbf{w}_{it} - \bar{\mathbf{w}})\|^2 \right]^{1/2} \left[ E \|\mathbf{g}_t\|^2 \right]^{1/2} < K,$$

and

$$E \left\| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i [(\mathbf{w}_{it} - \bar{\mathbf{w}}) \mathbf{g}_t' + \mathbf{g}_t (\mathbf{w}_{it} - \bar{\mathbf{w}})'] \right\| \leq \frac{2K}{N} \sum_{i=1}^N E |\eta_i| = O(N^{\delta-1}).$$

Using (A.6) and the above result in (A.3) we obtain

$$\mathbf{Q}_{P,N} = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})' + O_p(N^{\delta-1}),$$

which establishes that under  $\delta < 1$  (for a fixed  $T$  and as  $N \rightarrow \infty$ )

$$\mathbf{Q}_{P,N} \rightarrow_p \mathbf{\Omega}_P = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E [(\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})'] > 0.$$

Consider now the second component of (3.7), and note from (A.5) that since by assumption  $\eta_i$ ,  $u_{it}$ , and  $\mathbf{g}_t$  are distributed independently, then

$$\begin{aligned} E \left\| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i \mathbf{g}_t u_{it} \right\| &\leq \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E |\eta_i| E \|\mathbf{g}_t\| E |u_{it}| \\ &\leq \frac{K}{N} \sum_{i=1}^N E |\eta_i| = O(N^{\delta-1}). \end{aligned}$$

Hence, in view of (3.8) and using the above results we have

$$\hat{\beta}_P = \beta + \Omega_P^{-1} \left[ \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\eta_i + u_{it}) \right] + O_p(N^{\delta-1}). \quad (\text{A.7})$$

Furthermore

$$\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) \eta_i = \frac{1}{N} \sum_{i=1}^N (\bar{\mathbf{w}}_i - \bar{\mathbf{w}}) \eta_i,$$

and since by Assumption 1,  $\eta_i$  and  $\bar{\mathbf{w}}_i - \bar{\mathbf{w}}$  are independently distributed and by Assumption 4,  $E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\| < K$ , then

$$E \left\| \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) \eta_i \right\| \leq \frac{1}{N} \sum_{i=1}^N E |\eta_i| E \|\bar{\mathbf{w}}_i - \bar{\mathbf{w}}\| \leq \frac{K}{N} \sum_{i=1}^N E |\eta_i| = O(N^{\delta-1}).$$

Therefore, (A.7) simplifies further to

$$\hat{\beta}_P = \beta + \Omega_P^{-1} \left[ \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) u_{it} \right] + O_p(N^{\delta-1}). \quad (\text{A.8})$$

Using this result and noting that under Assumptions 2 and 4,

$$\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) u_{it} \rightarrow_p \mathbf{0}. \quad (\text{A.9})$$

Combining (A.8) and (A.9) yields the desired proposition.

## A.2 Proof of Proposition 3.3

For this result, noting that  $\mathbf{x}_{it} - \bar{\mathbf{x}}_i = (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) + \eta_i (\mathbf{g}_t - \bar{\mathbf{g}})$ , and  $y_{it} - \bar{y}_i = \beta' (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) + (u_{it} - \bar{u}_i)$ , we have

$$\begin{aligned} \mathbf{Q}_{FE,N} &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (\mathbf{x}_{it} - \bar{\mathbf{x}}_i)' \\ &= \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)' + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{g}_t - \bar{\mathbf{g}})' \\ &\quad + \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \eta_i (\mathbf{g}_t - \bar{\mathbf{g}}) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)' + \left( \frac{1}{N} \sum_{i=1}^N \eta_i^2 \right) \left( \frac{1}{T} \sum_{t=1}^T (\mathbf{g}_t - \bar{\mathbf{g}}) (\mathbf{g}_t - \bar{\mathbf{g}})' \right), \end{aligned}$$

and

$$\begin{aligned} \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{x}_{it} - \bar{\mathbf{x}}_i) (u_{it} - \bar{u}_i) &= \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T [(\mathbf{w}_{it} - \bar{\mathbf{w}}_i) + \eta_i (\mathbf{g}_t - \bar{\mathbf{g}})] (u_{it} - \bar{u}_i) \\ &= \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (u_{it} - \bar{u}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) \\ &\quad + \frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T \eta_i (u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}}). \end{aligned}$$



Under Assumptions 1-4, using the above results, and following the same line of reasoning as in Section 3.1 we have (for a fixed  $T$  and as  $N \rightarrow \infty$ )

$$\begin{aligned}\mathbf{Q}_{FE,N} &= \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E [(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)'] \\ &\quad + \left( \frac{1}{N} \sum_{i=1}^N E(\eta_i^2) \right) \left( \frac{1}{T} \sum_{t=1}^T E[(\mathbf{g}_t - \bar{\mathbf{g}})(\mathbf{g}_t - \bar{\mathbf{g}})'] \right) \\ &= \mathbf{\Omega}_{FE} + O_p(N^{\delta-1}),\end{aligned}\tag{A.10}$$

where

$$\mathbf{\Omega}_{FE} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E [(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)'].\tag{A.11}$$

Similarly, since  $\eta_i$  is distributed independently of  $u_{it}$  and  $\mathbf{g}_t$ , then

$$\begin{aligned}E \left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \eta_i (u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}}) \right| &\leq \frac{1}{\sqrt{NT}} \left| \sum_{i=1}^N \sum_{t=1}^T E |\eta_i| E |(u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}})| \right| \\ &\leq \sup_i E |(u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}})| \left( N^{-1/2} \sum_{i=1}^N E |\eta_i| \right).\end{aligned}$$

But  $E |(u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}})| \leq [E (u_{it} - \bar{u}_i)^2]^{1/2} [E \|\mathbf{g}_t - \bar{\mathbf{g}}\|^2]^{1/2} < K$ , and by Assumptions 1 and 2, it follows that

$$E \left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T \eta_i (u_{it} - \bar{u}_i) (\mathbf{g}_t - \bar{\mathbf{g}}) \right| \leq O(N^{\delta-1/2}).$$

Finally, under Assumptions 2-4, using standard results from panel data literature we have

$$\frac{1}{T\sqrt{N}} \sum_{i=1}^N \sum_{t=1}^T (u_{it} - \bar{u}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) \rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \mathbf{\Omega}_{FE}),$$

where  $\mathbf{\Omega}_{FE}$  is already defined by (A.11).

To establish the relative asymptotic efficiency of  $\hat{\beta}_P$  we first note that

$$\left[ \text{AsyVar} \left( \sqrt{TN} \hat{\beta}_P \right) \right]^{-1} - \left[ \text{AsyVar} \left( \sqrt{TN} \hat{\beta}_{FE} \right) \right]^{-1} = \sigma_u^{-2} [\mathbf{\Omega}_P - \mathbf{\Omega}_{FE}].\tag{A.12}$$

Also, we note that since

$$\frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}) (\mathbf{w}_{it} - \bar{\mathbf{w}})' = \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T (\mathbf{w}_{it} - \bar{\mathbf{w}}_i) (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)' + \frac{1}{N} \sum_{i=1}^N (\bar{\mathbf{w}}_i - \bar{\mathbf{w}}) (\bar{\mathbf{w}}_i - \bar{\mathbf{w}})',$$

then

$$\mathbf{\Omega}_P = \mathbf{\Omega}_{FE} + \mathbf{\Omega}_C,\tag{A.13}$$

where

$$\mathbf{\Omega}_C = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N E [(\bar{\mathbf{w}}_i - \bar{\mathbf{w}}) (\bar{\mathbf{w}}_i - \bar{\mathbf{w}})'],$$

and by Assumption 5,  $\mathbf{\Omega}_C$  is a positive definite matrix. Using (A.13) in (A.12) we have

$$\left[ \text{AsyVar} \left( \sqrt{TN} \hat{\beta}_P \right) \right]^{-1} - \left[ \text{AsyVar} \left( \sqrt{TN} \hat{\beta}_{FE} \right) \right]^{-1} = \sigma_u^{-2} \mathbf{\Omega}_C > 0,$$

and hence

$$\text{AsyVar} \left( \sqrt{TN} \hat{\beta}_{FE} \right) > \text{AsyVar} \left( \sqrt{TN} \hat{\beta}_P \right),$$

which gives the desired results in the proposition.

### A.3 Derivation of equation (4.11)

To show this result, we notice that

$$\begin{aligned} \text{Var}(\sqrt{NT}\hat{\mathbf{q}}) &= NT \times \text{Var}(\hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE}) \\ &= NT\text{Var}(\hat{\boldsymbol{\beta}}_P) + NT\text{Var}(\hat{\boldsymbol{\beta}}_{FE}) - NT\text{Cov}(\hat{\boldsymbol{\beta}}_{FE}, \hat{\boldsymbol{\beta}}_P) - NT\text{Cov}(\hat{\boldsymbol{\beta}}_P, \hat{\boldsymbol{\beta}}_{FE}), \end{aligned}$$

and it can be shown that

$$\begin{aligned} NT \times \text{Var}(\hat{\boldsymbol{\beta}}_P - \hat{\boldsymbol{\beta}}_{FE}) &= \mathbf{Q}_P^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T E[(\mathbf{w}_{it} - \bar{\mathbf{w}})(\mathbf{w}_{it'} - \bar{\mathbf{w}}) u_{it} u_{it'}] \mathbf{Q}_P^{-1} \\ &\quad + \mathbf{Q}_{FE}^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E[(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it'} - \bar{\mathbf{w}}_i)' u_{it} u_{it'}] \mathbf{Q}_{FE}^{-1} \\ &\quad - \mathbf{Q}_{FE}^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E[(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it'} - \bar{\mathbf{w}})' u_{it} u_{it'}] \mathbf{Q}_P^{-1} \\ &\quad - \mathbf{Q}_P^{-1} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T E[(\mathbf{w}_{it} - \bar{\mathbf{w}})(\mathbf{w}_{it'} - \bar{\mathbf{w}}_i)' u_{it} u_{it'}] \mathbf{Q}_{FE}^{-1} + O(N^{\delta-1}). \quad (\text{A.14}) \end{aligned}$$

Then by letting

$$\mathbf{V}_P = \lim_{N \rightarrow \infty} \mathbf{V}_{P,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}})(\mathbf{w}_{it} - \bar{\mathbf{w}})',$$

with  $\gamma_i(t, t') = E(u_{it} u_{jt'} | \mathbf{X}) = \gamma_i(t, t')$ , if  $i = j$  and  $t \neq t'$ , and

$$\mathbf{V}_{FE} = \lim_{N \rightarrow \infty} \mathbf{V}_{FE,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it} - \bar{\mathbf{w}}_i)',$$

and

$$\mathbf{V}_{FEP} = \lim_{N \rightarrow \infty} \mathbf{V}_{FEP,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it} - \bar{\mathbf{w}}_i)(\mathbf{w}_{it'} - \bar{\mathbf{w}})',$$

and

$$\mathbf{V}_{PFE} = \lim_{N \rightarrow \infty} \mathbf{V}_{PFE,N} = \lim_{N \rightarrow \infty} \frac{1}{NT} \sum_{i=1}^N \sum_{t=1}^T \sum_{t'=1}^T \gamma_i(t, t') (\mathbf{w}_{it'} - \bar{\mathbf{w}})(\mathbf{w}_{it} - \bar{\mathbf{w}})',$$

we can obtain the desired results.

### A.4 Proof of Proposition 5.1

The derivation is straightforward by following Wang et al. (2016). For (i), one can apply the Cramer-Wold device, since it is shown in the Proposition 3.3,

$$\begin{aligned} \sqrt{N}(\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}) &\rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \boldsymbol{\Omega}_P^{-1}), \\ \sqrt{N}(\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta}) &\rightarrow_d N(\mathbf{0}, \sigma_u^2 T^{-1} \boldsymbol{\Omega}_{FE}^{-1}), \end{aligned}$$

and it can be shown that under Assumptions 1-5,

$$\text{Cov}[\sqrt{N}(\hat{\boldsymbol{\beta}}_P - \boldsymbol{\beta}), \sqrt{N}(\hat{\boldsymbol{\beta}}_{FE} - \boldsymbol{\beta})] = \mathbf{V}_P = \sigma_u^2 T^{-1} \boldsymbol{\Omega}_P^{-1}. \quad (\text{A.15})$$

Result (ii) is established in the paper. To establish result (iii) we first note that under Assumption 1 and for  $\delta < \frac{1}{2}$  we have

$$\begin{aligned} Cov\left[\left(\hat{\beta}_P - \beta\right), \left(\hat{\beta}_{FE} - \hat{\beta}_P\right)\right] &= Cov\left(\hat{\beta}_P, \hat{\beta}_{FE}\right) - Cov\left(\hat{\beta}_P, \hat{\beta}_P\right) \\ &= Cov\left(\hat{\beta}_P, \hat{\beta}_P\right) - Cov\left(\hat{\beta}_P, \hat{\beta}_P\right) \\ &= 0, \end{aligned}$$

where the penultimate identity follows using (A.15). Then by continuous mapping theorem we obtain

$$\begin{aligned} \sqrt{N}\left(\hat{\beta}_{Pretest} - \beta\right) &= \sqrt{N}\left(\hat{\beta}_P - \beta\right) + 1\left(\tilde{H}_N > \chi_{k,1-\tau}^2\right)\sqrt{N}\left(\hat{\beta}_{FE} - \hat{\beta}_P\right) \\ &\rightarrow d\psi_1 + \mathbf{\Omega}^{-1}\psi_3 1\left(\varphi > \chi_{k,1-\tau}^2\right), \end{aligned}$$

as required.

## A.5 Additional simulation results

Here we provide additional simulation results for DGP (6.1) when  $T = 3$  and 10.

Table A1: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 3$  and  $0 \leq \delta \leq 0.44$

N		$\delta$																			
		0				0.25				0.4				0.42				0.44			
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest	
100	Estim	0.9997	1.0004	0.9997	0.9972	1.0120	1.0097		0.9994	1.0316	1.0216		1.0032	1.0422	1.0323		1.0027	1.0390	1.0293		
	Bias	-0.0003	0.0004	-0.0003	-0.0028	0.0120	0.0097		-0.0006	0.0316	0.0216		0.0032	0.0422	0.0323		0.0027	0.0390	0.0293		
	aBias	0.0723	0.0323	0.0382	0.0754	0.0299	0.0391		0.0762	0.0380	0.485		0.0710	0.0471	0.0529		0.0745	0.0437	0.0516		
	rmse	0.0895	0.0407	0.0527	0.0951	0.0373	0.0579		0.0957	0.0472	0.0676		0.0889	0.0573	0.0675		0.0936	0.0532	0.0687		
500	Estim	1.0019	1.0001	1.0003	0.9973	1.0048	1.0038		1.0013	1.0154	1.0122		1.0001	1.0189	1.0142		0.9992	1.0142	1.0112		
	Bias	0.0019	0.0001	0.0003	-0.0027	0.0048	0.0038		0.0013	0.0154	0.0122		0.0001	0.0189	0.0142		-0.0008	0.0142	0.0112		
	aBias	0.0338	0.0132	0.0165	0.0326	0.0132	0.0165		0.0324	0.0187	0.0219		0.0306	0.0204	0.0227		0.0323	0.0174	0.0207		
	rmse	0.0423	0.0166	0.0251	0.0406	0.0166	0.0242		0.0408	0.0229	0.0293		0.0381	0.0240	0.0285		0.0402	0.0213	0.0277		
1000	Estim	1.0001	1.0005	1.0004	1.0012	1.0045	1.0045		0.9999	1.0090	1.0074		1.0005	1.0106	1.0083		0.9999	1.0125	1.0099		
	Bias	0.0001	0.0005	0.0004	0.0012	0.0045	0.0045		-0.0001	0.0090	0.0074		0.0005	0.0106	0.0083		-0.0001	0.0125	0.0099		
	aBias	0.0236	0.0092	0.0114	0.0243	0.0102	0.0127		0.0232	0.0124	0.0141		0.0232	0.0130	0.0148		0.0233	0.0141	0.0159		
	rmse	0.0295	0.0117	0.0173	0.0308	0.0126	0.0187		0.0289	0.0141	0.0185		0.0289	0.0157	0.0193		0.0288	0.0168	0.0200		
2000	Estim	1.0003	1.0005	1.0004	0.9997	1.0026	1.0021		1.0008	1.0037	1.0032		0.9994	1.0050	1.0038		0.9997	1.0081	1.0056		
	Bias	0.0003	0.0005	0.0004	-0.0003	0.0026	0.0021		0.0008	0.0037	0.0032		-0.0006	0.0050	0.0038		-0.0003	0.0081	0.0056		
	aBias	0.0165	0.0064	0.0078	0.0166	0.0067	0.0086		0.0160	0.0069	0.0083		0.0167	0.0078	0.0094		0.0156	0.0094	0.0113		
	rmse	0.0206	0.0080	0.0114	0.0209	0.0084	0.0131		0.0203	0.0087	0.0122		0.0209	0.0098	0.0132		0.0197	0.0112	0.0148		

See notes of Table 1.

Table A2: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 3$  and  $0.46 \leq \delta \leq 1$

		$\delta$																				
		0.46				0.48				0.5				0.75				1				
$N$		FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest
100	Estim	1.0027	1.0390	1.0293	1.0058	1.0462	1.0357	1.0054	1.0597	1.0423	1.0054	1.0597	1.0423	0.9981	1.1099	1.0420	0.9951	1.1361	0.9983	0.9951	1.1361	0.9983
	Bias	0.0027	0.0390	0.0293	0.0058	0.0462	0.0357	0.0054	0.0597	0.0423	0.0054	0.0597	0.0423	-0.0019	0.1099	0.0420	-0.0049	0.1361	-0.0017	-0.0049	0.1361	-0.0017
	aBias	0.0745	0.0437	0.0516	0.0760	0.0504	0.0585	0.0757	0.0619	0.0677	0.0757	0.0619	0.0677	0.0657	0.1100	0.0931	0.0668	0.1361	0.0693	0.0668	0.1361	0.0693
	rmse	0.0936	0.0532	0.0687	0.0944	0.0588	0.0732	0.0949	0.0709	0.0801	0.0949	0.0709	0.0801	0.0841	0.1160	0.1056	0.0831	0.1403	0.0858	0.0831	0.1403	0.0858
500	Estim	1.0013	1.0132	1.0100	0.9998	1.0163	1.0124	1.0007	1.0222	1.0167	1.0007	1.0222	1.0167	0.9975	1.0708	1.0098	1.0017	1.1436	1.0017	1.0017	1.1436	1.0017
	Bias	0.0013	0.0132	0.0100	-0.0002	0.0163	0.0124	0.0007	0.0222	0.0167	0.0007	0.0222	0.0167	-0.0025	0.0708	0.0098	0.0017	0.1436	0.0017	0.0017	0.1436	0.0017
	aBias	0.0323	0.0167	0.0200	0.0318	0.0187	0.0226	0.0318	0.0234	0.0260	0.0318	0.0234	0.0260	0.0301	0.0708	0.0415	0.0279	0.1436	0.0279	0.0279	0.1436	0.0279
	rmse	0.0409	0.0206	0.0273	0.0398	0.0227	0.0299	0.0397	0.0275	0.0318	0.0397	0.0275	0.0318	0.0379	0.0726	0.0498	0.0350	0.1444	0.0350	0.0350	0.1444	0.0350
1000	Estim	1.0006	1.0116	1.0093	0.9999	1.0120	1.0089	1.0004	1.0151	1.0115	1.0004	1.0151	1.0115	1.0015	1.0592	1.0066	0.9994	1.1428	0.9994	0.9994	1.1428	0.9994
	Bias	0.0006	0.0116	0.0093	-0.0001	0.0120	0.0089	0.0004	0.0151	0.0115	0.0004	0.0151	0.0115	0.0015	0.0592	0.0066	-0.0006	0.1428	-0.0004	-0.0006	0.1428	-0.0004
	aBias	0.0227	0.0135	0.0160	0.0221	0.0136	0.0157	0.0227	0.0163	0.0182	0.0227	0.0163	0.0182	0.0216	0.0592	0.0265	0.0196	0.1428	0.0196	0.0196	0.1428	0.0196
	rmse	0.0283	0.0163	0.0208	0.0281	0.0165	0.0206	0.0284	0.0193	0.0226	0.0284	0.0193	0.0226	0.0268	0.0602	0.0333	0.0248	0.1432	0.0248	0.0248	0.1432	0.0248
2000	Estim	1.0011	1.0084	1.0064	1.0003	1.0098	1.0082	0.9996	1.0095	1.0078	0.9996	1.0095	1.0078	1.0005	1.0473	1.0016	1.0009	1.1432	1.0009	1.0009	1.1432	1.0009
	Bias	0.0011	0.0084	0.0064	0.0003	0.0098	0.0082	-0.0004	0.0095	0.0078	-0.0004	0.0095	0.0078	0.0005	0.0473	0.0016	0.0009	0.1432	0.0009	0.0009	0.1432	0.0009
	aBias	0.0171	0.0095	0.0112	0.0155	0.0107	0.0122	0.0150	0.0105	0.0116	0.0150	0.0105	0.0116	0.0155	0.0473	0.0166	0.0142	0.1432	0.0142	0.0142	0.1432	0.0142
	rmse	0.0215	0.0115	0.0148	0.0197	0.0128	0.0156	0.0190	0.0126	0.0147	0.0190	0.0126	0.0147	0.0195	0.0480	0.0212	0.0175	0.1435	0.0175	0.0175	0.1435	0.0175

See notes of Table 1.

Table A3: Simulation results for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 3$  and  $0 \leq \delta \leq 0.44$

$N$		$\delta$																			
		0				0.25				0.4				0.42				0.44			
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest	
100	Estim	1.9956	2.0005	2.0008	1.9990	2.0060	2.0028	2.0001	2.0417	2.0289	1.9996	2.0383	2.0271	2.0049	2.0496	2.0000	2.0049	2.0496	2.0000	2.0496	2.0000
	Bias	-0.0044	0.0005	0.0008	-0.0010	0.0060	0.0028	0.0001	0.0417	0.0289	-0.0004	0.0383	0.0271	0.0049	0.0496	0.0000	0.0049	0.0496	0.0000	0.0496	0.0000
	aBias	0.0777	0.0284	0.0369	0.0755	0.0293	0.0373	0.0745	0.0446	0.0532	0.0734	0.0425	0.0510	0.0723	0.0513	0.0723	0.0425	0.0510	0.0723	0.0425	0.0510
	rmse	0.0970	0.0351	0.0566	0.0944	0.0365	0.0550	0.0929	0.0531	0.0679	0.0923	0.0504	0.0672	0.0908	0.0593	0.0908	0.0593	0.0593	0.0908	0.0593	0.0593
500	Estim	2.0002	2.0005	2.0002	2.0004	2.0103	2.0072	1.9978	2.0130	2.0093	2.0013	2.0157	2.0123	1.9992	2.0129	2.0000	2.0129	2.0129	2.0000	2.0129	2.0000
	Bias	0.0002	0.0005	0.0002	0.0004	0.0103	0.0072	-0.0022	0.0130	0.0093	0.0013	0.0157	0.0123	-0.0008	0.0129	0.0000	0.0129	0.0129	0.0000	0.0129	0.0000
	aBias	0.0348	0.0138	0.0177	0.0347	0.0155	0.0187	0.0344	0.0169	0.0207	0.0324	0.0182	0.0210	0.0335	0.0167	0.0335	0.0167	0.0167	0.0335	0.0167	0.0335
	rmse	0.0434	0.0171	0.0265	0.0431	0.0193	0.0269	0.0426	0.0205	0.0287	0.0410	0.0221	0.0276	0.0416	0.0204	0.0416	0.0204	0.0204	0.0416	0.0204	0.0416
1000	Estim	2.0009	2.0000	1.9997	2.0006	2.0038	2.0034	2.0007	2.0076	2.0066	2.0007	2.0076	2.0063	2.0013	2.0124	2.0000	2.0124	2.0124	2.0000	2.0124	2.0000
	Bias	0.0009	0.0000	-0.0003	0.0006	0.0038	0.0034	0.0007	0.0076	0.0066	0.0007	0.0076	0.0063	0.0013	0.0124	0.0000	0.0124	0.0124	0.0000	0.0124	0.0000
	aBias	0.0243	0.0093	0.0116	0.0240	0.0240	0.0120	0.0245	0.0113	0.0138	0.0232	0.0113	0.0137	0.0246	0.0140	0.0246	0.0140	0.0140	0.0246	0.0140	0.0246
	rmse	0.0304	0.0116	0.0179	0.0304	0.0304	0.0183	0.0307	0.0140	0.0195	0.0289	0.0138	0.0195	0.0310	0.0170	0.0310	0.0170	0.0170	0.0310	0.0170	0.0310
2000	Estim	1.9994	1.9996	1.9994	1.9994	2.0024	2.0018	1.9992	2.0050	2.0039	2.0003	2.0047	2.0037	1.9997	2.0087	2.0000	2.0087	2.0087	2.0000	2.0087	2.0000
	Bias	-0.0006	-0.0004	-0.0006	-0.0006	0.0024	0.0018	-0.0008	0.0050	0.0039	0.0003	0.0047	0.0037	-0.0003	0.0087	0.0000	0.0087	0.0087	0.0000	0.0087	0.0000
	aBias	0.0170	0.0066	0.0083	0.0164	0.0069	0.0086	0.0166	0.0078	0.0091	0.0157	0.0075	0.0087	0.0160	0.0097	0.0160	0.0097	0.0097	0.0160	0.0097	0.0160
	rmse	0.0216	0.0082	0.0124	0.0207	0.0086	0.0129	0.0207	0.0096	0.0129	0.0199	0.0092	0.0121	0.0204	0.0116	0.0204	0.0116	0.0116	0.0204	0.0116	0.0204

See notes of Table 1.

Table A4: Simulation results for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 3$  and  $0.46 \leq \delta \leq 1$

		$\delta$																				
		0.46				0.48				0.5				0.75				1				
$N$		FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest
100	Estim	2.0049	2.0496	2.0356	1.9999	2.0472	2.0315	1.9999	2.0482	2.0355	2.0013	2.0822	2.0336	2.0000	2.1617	2.0336	2.0000	2.1617	2.0336	2.0000	2.1617	2.0336
	Bias	0.0049	0.0496	0.0356	-0.0001	0.0472	0.0315	-0.0001	0.0482	0.0355	0.0013	0.0822	0.0336	0.0000	0.1617	0.0336	0.0000	0.1617	0.0336	0.0000	0.1617	0.0336
	aBias	0.0723	0.0513	0.0593	0.0759	0.0500	0.0600	0.0660	0.0499	0.0572	0.0657	0.0825	0.0813	0.0603	0.1617	0.0813	0.0603	0.1617	0.0813	0.0603	0.1617	0.0813
	rmse	0.0908	0.0593	0.0727	0.0954	0.0582	0.0751	0.0834	0.0585	0.0707	0.0841	0.0894	0.0921	0.0757	0.1655	0.0921	0.0757	0.1655	0.0921	0.0757	0.1655	0.0921
500	Estim	2.0001	2.0130	2.0106	1.9989	2.0198	2.0153	1.9991	2.0195	2.0152	1.9995	2.0678	2.0109	1.9984	2.1376	2.0109	1.9984	2.1376	2.0109	1.9984	2.1376	2.0109
	Bias	0.0001	0.0130	0.0106	-0.0011	0.0198	0.0153	-0.0009	0.0195	0.0152	-0.0005	0.0678	0.0109	-0.0016	0.1376	0.0109	-0.0016	0.1376	0.0109	-0.0016	0.1376	0.0109
	aBias	0.0345	0.0166	0.0206	0.0328	0.0215	0.0249	0.0319	0.0214	0.0244	0.0314	0.0678	0.0410	0.0265	0.1376	0.0410	0.0265	0.1376	0.0410	0.0265	0.1376	0.0410
	rmse	0.0439	0.0205	0.0292	0.0416	0.0255	0.0320	0.0397	0.0244	0.0307	0.0393	0.0698	0.0492	0.0332	0.1385	0.0492	0.0332	0.1385	0.0492	0.0332	0.1385	0.0492
1000	Estim	1.9998	2.0159	2.0114	1.9993	2.0150	2.0110	1.9990	2.0145	2.0111	1.9982	2.0560	2.0036	2.0001	2.1390	2.0036	2.0001	2.1390	2.0036	2.0001	2.1390	2.0036
	Bias	-0.0002	0.0159	0.0114	-0.0007	0.0150	0.0110	-0.0010	0.0145	0.0111	-0.0018	0.0560	0.0036	0.0001	0.1390	0.0036	0.0001	0.1390	0.0036	0.0001	0.1390	0.0036
	aBias	0.0244	0.0167	0.0191	0.0241	0.0160	0.0188	0.0235	0.0156	0.0183	0.0217	0.0560	0.0269	0.0192	0.1390	0.0269	0.0192	0.1390	0.0269	0.0192	0.1390	0.0269
	rmse	0.0303	0.0195	0.0238	0.0306	0.0187	0.0237	0.0300	0.0185	0.0235	0.0270	0.0570	0.0335	0.0241	0.1394	0.0335	0.0241	0.1394	0.0335	0.0241	0.1394	0.0335
2000	Estim	2.0000	2.0102	2.0077	1.9996	2.0097	2.0078	1.9999	2.0094	2.0073	1.9993	2.0482	2.0003	1.9992	2.1389	2.0003	1.9992	2.1389	2.0003	1.9992	2.1389	2.0003
	Bias	0.0000	0.0102	0.0077	-0.0004	0.0097	0.0078	-0.0001	0.0094	0.0073	-0.0007	0.0482	0.0003	-0.0008	0.1389	0.0003	-0.0008	0.1389	0.0003	-0.0008	0.1389	0.0003
	aBias	0.0169	0.0109	0.0126	0.0162	0.0108	0.0123	0.0168	0.0104	0.0120	0.0155	0.0482	0.0166	0.0139	0.1389	0.0166	0.0139	0.1389	0.0166	0.0139	0.1389	0.0166
	rmse	0.0212	0.0129	0.0160	0.0203	0.0128	0.0156	0.0211	0.0123	0.0155	0.0195	0.0488	0.0213	0.0173	0.1391	0.0213	0.0173	0.1391	0.0213	0.0173	0.1391	0.0213

See notes of Table 1.

Table A5: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 10$  and  $0 \leq \delta \leq 0.44$

N		$\delta$																			
		0				0.25				0.4				0.42				0.44			
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest	
100	Estim	0.9998	0.9999	1.0006		0.9996	1.0319	1.0283		0.9999	1.0418	1.0198		0.9996	1.0470	1.0157		0.9999	1.0453	1.012	
	Bias	-0.0002	-0.0001	0.0006		-0.0004	0.0319	0.0283		-0.0001	0.0418	0.0198		-0.0004	0.0470	0.0157		-0.0001	0.0453	0.012	
	aBias	0.0291	0.0173	0.0196		0.0259	0.0324	0.0332		0.0261	0.0418	0.0375		0.0272	0.0471	0.0367		0.0261	0.0453	0.034	
	rmse	0.0368	0.0216	0.0263		0.0327	0.0367	0.0378		0.0328	0.0452	0.0428		0.0338	0.0506	0.0425		0.0332	0.0485	0.040	
500	Estim	0.9989	0.9997	0.9993		1.0000	1.0136	1.0111		0.9996	1.0253	1.0116		0.9998	1.0290	1.0080		1.0006	1.0273	1.0099	
	Bias	-0.0011	-0.0003	-0.0007		0.0000	0.0136	0.0111		-0.0004	0.0253	0.0116		-0.0002	0.0290	0.0080		0.0006	0.0273	0.0099	
	aBias	0.0133	0.0072	0.0080		0.0124	0.0136	0.0146		0.0126	0.0253	0.0207		0.0116	0.0290	0.0185		0.0129	0.0273	0.020	
	rmse	0.0166	0.0090	0.0107		0.0154	0.0140	0.0170		0.0157	0.0267	0.0233		0.0145	0.0300	0.0200		0.0158	0.0284	0.023	
1000	Estim	1.0000	0.9999	1.0000		0.9996	1.0101	1.0076		1.0002	1.0179	1.0081		0.9999	1.0210	1.0076		0.9999	1.0219	1.005	
	Bias	0.0000	-0.0001	0.0000		-0.0004	0.0101	0.0076		0.0002	0.0179	0.0081		-0.0001	0.0210	0.0076		-0.0001	0.0219	0.005	
	aBias	0.0094	0.0049	0.0055		0.0096	0.0103	0.0109		0.0086	0.0179	0.0142		0.0088	0.0210	0.0149		0.0088	0.0219	0.012	
	rmse	0.0120	0.0055	0.0076		0.0119	0.0118	0.0127		0.0108	0.0188	0.0160		0.0110	0.0209	0.0170		0.0110	0.0227	0.022	
2000	Estim	0.9998	1.0001	0.9999		1.0005	1.0067	1.0049		1.0002	1.0113	1.0047		1.0002	1.0134	1.0043		1.0000	1.0163	1.001	
	Bias	-0.0002	0.0001	-0.0001		0.0005	0.0067	0.0049		0.0002	0.0113	0.0047		0.0002	0.0134	0.0043		0.0000	0.0163	0.001	
	aBias	0.0067	0.0034	0.0040		0.0066	0.0068	0.0072		0.0065	0.0113	0.0094		0.0065	0.0134	0.0098		0.0062	0.0163	0.007	
	rmse	0.0083	0.0040	0.0053		0.0082	0.0078	0.0084		0.0081	0.0121	0.0107		0.0082	0.0141	0.0113		0.0078	0.0168	0.009	

See notes of Table 1.



Table A6: Simulation results for  $\beta_1$  ( $\beta_1 = 1$ ) when  $T = 10$  and  $0.46 \leq \delta \leq 1$

$N$		$\delta$																			
		0.46				0.48				0.5				0.75				1			
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest	
100	Estim	0.9999	1.0453	1.0121	0.9978	1.0498	1.0103		0.9993	1.0531	1.0098		0.9999	1.0684	1.0000		0.9992	1.0622	0.9992		
	Bias	-0.0001	0.0453	0.0121	-0.0022	0.0498	0.0103		-0.0007	0.0531	0.0098		-0.0001	0.0684	0.0000		-0.0008	0.0622	-0.0008		
	aBias	0.0261	0.0453	0.0342	0.0264	0.0498	0.0353		0.0280	0.0531	0.0363		0.0239	0.0684	0.0239		0.0232	0.0622	0.0232		
	rmse	0.0332	0.0485	0.0403	0.0337	0.0530	0.0421		0.0349	0.0557	0.0431		0.0297	0.0704	0.0297		0.0291	0.0642	0.0291		
500	Estim	1.0009	1.0278	1.0085	0.9996	1.0301	1.0050		0.9997	1.0350	1.0022		0.9996	1.0581	0.9996		0.9998	1.0678	0.9998		
	Bias	0.0009	0.0278	0.0085	-0.0004	0.0301	0.0050		-0.0003	0.0350	0.0022		-0.0004	0.0581	-0.0004		-0.0002	0.0678	-0.0002		
	aBias	0.0115	0.0278	0.0179	0.0120	0.0301	0.0167		0.0117	0.0350	0.0141		0.0104	0.0581	0.0104		0.0099	0.0678	0.0099		
	rmse	0.0146	0.0289	0.0213	0.0151	0.0313	0.0206		0.0145	0.0358	0.0180		0.0130	0.0586	0.0130		0.0125	0.0681	0.0125		
1000	Estim	0.9996	1.0229	1.0023	0.9988	1.0246	0.9998		0.9992	1.0278	1.0001		0.9998	1.0543	0.9998		0.9999	1.0680	0.9999		
	Bias	-0.0004	0.0229	0.0023	-0.0012	0.0246	-0.0002		-0.0008	0.0278	0.0001		-0.0002	0.0543	-0.0002		-0.0001	0.0680	-0.0001		
	aBias	0.0090	0.0230	0.0114	0.0086	0.0246	0.0098		0.0086	0.0278	0.0094		0.0079	0.0543	0.0079		0.0070	0.0680	0.0070		
	rmse	0.0114	0.0238	0.0144	0.0108	0.0253	0.0123		0.0108	0.0285	0.0122		0.0098	0.0546	0.0098		0.0087	0.0682	0.0087		
2000	Estim	1.0000	1.0178	1.0016	1.0001	1.0193	1.0009		1.0003	1.0196	1.0012		0.9998	1.0501	0.9998		1.0001	1.0679	1.0001		
	Bias	0.0000	0.0178	0.0016	0.0001	0.0193	0.0009		0.0003	0.0196	0.0012		-0.0002	0.0501	-0.0002		0.0001	0.0679	0.0001		
	aBias	0.0063	0.0178	0.0078	0.0064	0.0193	0.0072		0.0065	0.0196	0.0074		0.0056	0.0501	0.0056		0.0050	0.0679	0.0050		
	rmse	0.0.079	0.0182	0.0099	0.0079	0.0197	0.0092		0.0080	0.0200	0.0094		0.0069	0.0502	0.0069		0.0063	0.0680	0.0063		

See notes of Table 1.

Table A7: Simulation results for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 10$  and  $0 \leq \delta \leq 0.44$

$N$		$\delta$																				
		0				0.25				0.4				0.42				0.44				
		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		FE	PLS	Pretest		
100	Estim	2.0016	2.0004	2.0006	2.0012	2.0223	2.0208	2.0009	2.0435	2.0209	2.0007	2.0433	2.0162	2.0004	2.0470	2.0130	2.0004	2.0470	2.0130	2.0004	2.0470	2.0130
	Bias	0.0016	0.0004	0.0006	0.0012	0.0223	0.0288	0.0009	0.0435	0.0209	0.0007	0.0433	0.0162	0.0004	0.0470	0.0130	0.0004	0.0470	0.0130	0.0004	0.0470	0.0130
	aBias	0.0.312	0.0156	0.0189	0.0278	0.0244	0.0265	0.0272	0.0435	0.0388	0.0264	0.0433	0.0365	0.0261	0.0470	0.0355	0.0261	0.0470	0.0355	0.0261	0.0470	0.0355
	rmse	0.0393	0.0201	0.0270	0.0352	0.0288	0.0321	0.0344	0.0465	0.0438	0.0338	0.0461	0.0422	0.0327	0.0496	0.0408	0.0327	0.0496	0.0408	0.0327	0.0496	0.0408
500	Estim	1.9995	1.9998	1.9998	2.0005	2.0177	2.0146	1.9995	2.0239	2.0105	2.0006	2.0267	2.0077	1.9999	2.0264	2.0089	1.9999	2.0264	2.0089	1.9999	2.0264	2.0089
	Bias	-0.0005	-0.0002	-0.0002	0.0005	0.0177	0.0146	-0.0005	0.0.239	0.0105	0.0006	0.0267	0.0077	-0.0001	0.0264	0.0089	-0.0001	0.0264	0.0089	-0.0001	0.0264	0.0089
	aBias	0.0137	0.0075	0.0084	0.0134	0.0177	0.0180	0.0130	0.0239	0.0105	0.0126	0.0267	0.0184	0.0124	0.0265	0.0199	0.0124	0.0265	0.0199	0.0124	0.0265	0.0199
	rmse	0.0172	0.0094	0.0112	0.0168	0.0178	0.0201	0.0162	0.0253	0.0227	0.0158	0.0279	0.0216	0.0156	0.0277	0.0222	0.0156	0.0277	0.0222	0.0156	0.0277	0.0222
1000	Estim	2.0000	1.9997	1.9998	1.9999	2.0091	2.0072	2.0003	2.0169	2.0077	2.0004	2.0180	2.0068	2.0006	2.0220	2.0055	2.0006	2.0220	2.0055	2.0006	2.0220	2.0055
	Bias	0.0000	-0.0003	-0.0002	-0.0001	0.0091	0.0072	0.0003	0.0169	0.0077	0.0004	0.0180	0.0068	0.0006	0.0220	0.0055	0.0006	0.0220	0.0055	0.0006	0.0220	0.0055
	aBias	0.0096	0.0053	0.0061	0.0093	0.0094	0.0100	0.0092	0.0169	0.0141	0.0091	0.0180	0.0137	0.0093	0.0220	0.0139	0.0093	0.0220	0.0139	0.0093	0.0220	0.0139
	rmse	0.0122	0.0061	0.0086	0.0117	0.0109	0.0118	0.0115	0.0179	0.0159	0.0114	0.0190	0.0158	0.0116	0.0228	0.0163	0.0116	0.0228	0.0163	0.0116	0.0228	0.0163
2000	Estim	2.0001	1.9999	1.9999	1.9999	2.0064	2.0045	2.0001	2.0128	2.0051	2.0000	2.0136	2.0041	1.9997	2.0170	2.0019	1.9997	2.0170	2.0019	1.9997	2.0170	2.0019
	Bias	0.0001	-0.0001	-0.0001	-0.0001	0.0064	0.0045	0.0001	0.0128	0.0051	0.0000	0.0136	0.0041	-0.0003	0.0170	0.0018	-0.0003	0.0170	0.0018	-0.0003	0.0170	0.0018
	aBias	0.0068	0.0035	0.0040	0.0065	0.0066	0.0072	0.0063	0.0128	0.0098	0.0061	0.0136	0.0094	0.0064	0.0170	0.0089	0.0064	0.0170	0.0089	0.0064	0.0170	0.0089
	rmse	0.0086	0.0040	0.0056	0.0083	0.0077	0.0086	0.0079	0.0134	0.0112	0.0077	0.0142	0.0110	0.0079	0.0175	0.0109	0.0079	0.0175	0.0109	0.0079	0.0175	0.0109

See notes of Table 1.

Table A8: Simulation results for  $\beta_2$  ( $\beta_2 = 2$ ) when  $T = 10$  and  $0.46 \leq \delta \leq 1$

		$\delta$																				
		0.46				0.48				0.5				0.75				1				
$N$		FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest	FE	PLS	Pretest
100	Estim	2.0004	2.0470	2.0136	2.0008	2.0470	2.0125	2.0003	2.0455	2.0090	2.0010	2.0531	2.0010	2.0009	2.0740	2.0009	2.0009	2.0740	2.0009	2.0009	2.0740	2.0009
	Bias	0.0004	0.0470	0.0136	0.0008	0.0470	0.0125	0.0003	0.0455	0.0090	0.0010	0.0531	0.0010	0.0009	0.0740	0.0009	0.0009	0.0740	0.0009	0.0009	0.0740	0.0009
	aBias	0.0261	0.0470	0.0350	0.0258	0.0470	0.0340	0.0252	0.0455	0.0312	0.0235	0.0531	0.0235	0.0234	0.0740	0.0234	0.0234	0.0740	0.0234	0.0234	0.0740	0.0234
	rmse	0.0327	0.0496	0.0408	0.0325	0.0499	0.0400	0.0320	0.0484	0.0375	0.0295	0.0534	0.0295	0.0292	0.0759	0.0292	0.0292	0.0759	0.0292	0.0292	0.0759	0.0292
500	Estim	1.9998	2.0277	2.0079	2.0001	2.0327	2.0054	2.0006	2.0338	2.0030	1.9998	2.0560	1.9998	2.0003	2.0661	2.0003	2.0003	2.0661	2.0003	2.0003	2.0661	2.0003
	Bias	-0.0002	0.0277	0.0079	0.0001	0.0327	0.0054	0.0006	0.0338	0.0030	-0.0002	0.0560	-0.0002	0.0003	0.0661	0.0003	0.0003	0.0661	0.0003	0.0003	0.0661	0.0003
	aBias	0.0122	0.0277	0.0188	0.0131	0.0327	0.0178	0.0121	0.0338	0.0144	0.0106	0.0560	0.0106	0.0097	0.0661	0.0097	0.0097	0.0661	0.0097	0.0097	0.0661	0.0097
	rmse	0.0154	0.0288	0.0219	0.0162	0.0337	0.0218	0.0151	0.0347	0.0183	0.0133	0.0565	0.0133	0.0122	0.0665	0.0122	0.0122	0.0665	0.0122	0.0122	0.0665	0.0122
1000	Estim	1.9999	2.0260	2.0027	1.9998	2.0275	2.0009	1.9997	2.0276	2.0006	2.0002	2.0524	2.0002	2.0001	2.0656	2.0001	2.0001	2.0656	2.0001	2.0001	2.0656	2.0001
	Bias	-0.0001	0.0260	0.0027	-0.0002	0.0275	0.0009	-0.0003	0.0276	0.0006	0.0002	0.0524	0.0002	0.0001	0.0656	0.0001	0.0001	0.0656	0.0001	0.0001	0.0656	0.0001
	aBias	0.0095	0.0260	0.0121	0.0086	0.0275	0.0095	0.0089	0.0276	0.0098	0.0082	0.0524	0.0082	0.0071	0.0656	0.0071	0.0071	0.0656	0.0071	0.0071	0.0656	0.0071
	rmse	0.0118	0.0266	0.0254	0.0109	0.0281	0.0124	0.0112	0.0283	0.0126	0.0100	0.0527	0.0100	0.0089	0.0658	0.0089	0.0089	0.0658	0.0089	0.0089	0.0658	0.0089
2000	Estim	1.9999	2.0193	2.0015	1.9998	2.0197	2.0008	1.9997	2.0199	2.0005	1.9999	2.0507	1.9999	1.9998	2.0664	1.9998	1.9998	2.0664	1.9998	1.9998	2.0664	1.9998
	Bias	-0.0001	0.0193	0.0015	-0.0002	0.0917	0.0008	-0.0003	0.0199	0.0005	-0.0001	0.0507	-0.0001	-0.0002	0.0664	-0.0002	-0.0002	0.0664	-0.0002	-0.0002	0.0664	-0.0002
	aBias	0.0065	0.0193	0.0080	0.0064	0.0197	0.0073	0.0063	0.0199	0.0071	0.0054	0.0507	0.0054	0.0049	0.0664	0.0049	0.0049	0.0664	0.0049	0.0049	0.0664	0.0049
	rmse	0.0081	0.0198	0.0103	0.0080	0.0202	0.0094	0.0078	0.0203	0.0092	0.0067	0.0508	0.0067	0.0062	0.0665	0.0062	0.0062	0.0665	0.0062	0.0062	0.0665	0.0062

See notes of Table 1.